

Table 10.2 Vaccinations by source of information

Percentage of children age 12-23 months and children age 24-35 months who received specific vaccines at any time before the survey, by source of information (vaccination card or mother's report), and percentage vaccinated by appropriate age, [country year]

Vaccine	Children age 12-23 months:				Children age 24-35 months:			
	Vaccinated at any time before the survey according to:			Vaccinated by appropriate age ^{1,2}	Vaccinated at any time before the survey according to:			Vaccinated by appropriate age ^{1,2}
	Vaccination card	Mother's report	Either source		Vaccination card	Mother's report	Either source	
BCG								
HepB (birth dose)								
DPT-HepB-Hib³								
1								
2								
3								
Polio⁴								
0 (birth dose)								
1								
2								
3								
Pneumococcal⁵								
1								
2								
3								
Rotavirus⁵								
1								
2								
3								
[Measles containing vaccine]⁶								
1								
2		na	na	na		na	na	
All age appropriate vaccinations⁷								
No vaccinations								
Number of children								

na = Not applicable

¹For children age 12-23 months, by age 12 months; for children age 23-35 months, by age 12 months for all vaccines except [measles containing vaccine 2] which should be received by age 24 months

² For children whose information is based on the mother's report, the proportions of vaccinations given during the first and second years of life are assumed to be the same as for children with a written record of vaccination

³DPT-HepB-Hib is sometimes referred to as pentavalent [ADAPT TO FOLLOW NATIONAL IMMUNIZATION SCHEDULE. IF DPT, HEP B AND HIB ARE GIVEN SEPARATELY, PROVIDE COLUMNS FOR THE RECOMMENDED NUMBER OF DOSES OF EACH.]

⁴ Polio 0 is the polio vaccination given at birth. [DELETE IN COUNTRIES WHERE POLIO 0 (POLIO AT BIRTH) IS NOT PART OF THE IMMUNIZATION SCHEDULE.]

⁵[IF VACCINATION SCHEDULE ONLY USES TWO DOSES OF VACCINE, REMOVE THIRD ENTRY.]

⁶[ADAPT QUESTION LOCALLY TO USE THE NAME OF THE MEASES CONTAINING VACCINATION (MCV) USED INT THE COUNTRY: MEASES, MMR, OR MR.]

⁷ BCG, hepatitis B (birth dose), three doses of DPT-HepB-Hib, [three/two] doses of pneumococcal vaccine, [three/two] doses of rotavirus vaccine, [three/four] doses of polio vaccine, and one dose of measles containing vaccine (if 12-23 months) and two doses of measles containing vaccine (if 24-35 months)

Table 10.2 Vaccinations by source of information

Percentage of children age 12-23 months, children age 24-35 months, and children age 36-47 months who received specific vaccines at any time before the survey, by source of information (vaccination card or mother's report), and percentage vaccinated by appropriate age, [country year]

Vaccine	Children age 12-23 months:				Children age 24-35 months:				Children age 36-47 months:				
	Vaccinated at any time before the survey according to:			Vaccinated by appropriate age ^{1,2}	Vaccinated at any time before the survey according to:			Vaccinated by appropriate age ^{1,2}	Vaccinated at any time before the survey according to:			Vaccinated by appropriate age ^{1,2}	
	Vaccination card	Mother's report	Either source		Vaccination card	Mother's report	Either source		Vaccination card	Mother's report	Either source		
BCG													
HepB (birth dose)													
DPT-HepB-Hib³													
1													
2													
3													
Polio⁴													
0 (birth dose)													
1													
2													
3													
Pneumococcal⁵													
1													
2													
3													
Rotavirus⁵													
1													
2													
3													
[Measles containing vaccine]⁶													
1		na	na	na	na								
2		na	na	na	na	na	na	na	na	na			

All age appropriate vaccinations⁷

No vaccinations

Number of children

na = Not applicable

¹For children age 12-23 months, by age 12 months for all vaccines except [measles containing vaccine 1 and measles containing vaccine 2] which should be received by age 24 months and age 36 months, respectively. For children age 24-36 months, by age 24 months for all vaccines except [measles containing vaccine 2] which should be received by age 36 months; for children age 36-47 months, by age 36 months.

² For children whose information is based on the mother's report, the proportions of vaccinations given during the first and second years of life are assumed to be the same as for children with a written record of vaccination

³ DPT-HepB-Hib is sometimes referred to as pentavalent [ADAPT TO FOLLOW NATIONAL IMMUNIZATION SCHEDULE. IF DPT, HEP B AND HIB ARE GIVEN SEPARATELY, PROVIDE COLUMNS FOR THE RECOMMENDED NUMBER OF DOSES OF EACH.]

⁴ Polio 0 is the polio vaccination given at birth. [DELETE IN COUNTRIES WHERE POLIO 0 (POLIO AT BIRTH) IS NOT PART OF THE IMMUNIZATION SCHEDULE.]

⁵[IF VACCINATION SCHEDULE ONLY USES TWO DOSES OF VACCINE, REMOVE THIRD ENTRY.]

⁶[ADAPT QUESTION LOCALLY TO USE THE NAME OF THE MEASLES CONTAINING VACCINATION (MCV) USED IN THE COUNTRY: MEASLES, MMR, OR MR.]

⁷ BCG, hepatitis B (birth dose), three doses of DPT-HepB-Hib, [three/two] doses of pneumococcal vaccine, [three/two] doses of rotavirus vaccine, [three/four] doses of polio vaccine, and one dose of measles containing vaccine (if 24-36 months) and two doses of measles containing vaccine (if 36-47 months)

