

DHS 2015 CHILDREN DB

```
. use "\\ies\d5\foodsec\Share\SaraAC\1. raw DATABASES RWANDA\2015 DHS\download_July2018\RWKR70D\RWKR70FL.DTA"
. tab hw1 if hw1>=6 & hw1<=23
```

child's age in months	Freq.	Percent	Cum.
6	70	5.98	5.98
7	75	6.40	12.38
8	65	5.55	17.93
9	88	7.51	25.45
10	63	5.38	30.83
11	63	5.38	36.21
12	65	5.55	41.76
13	63	5.38	47.14
14	68	5.81	52.95
15	61	5.21	58.16
16	67	5.72	63.88
17	68	5.81	69.68
18	54	4.61	74.30
19	63	5.38	79.68
20	64	5.47	85.14
21	52	4.44	89.58
22	66	5.64	95.22
23	56	4.78	100.00
Total	1,171	100.00	

DHS 2012 CHILDREN DB

```
. use "\\ies\d5\foodsec\Share\SaraAC\1. raw DATABASES RWANDA\2010 DHS\RWKR61FL.DTA"
. tab hw1 if hw1>=6 & hw1<=23
```

child's age in months	Freq.	Percent	Cum.
6	58	4.78	4.78
7	74	6.10	10.87
8	72	5.93	16.80
9	70	5.77	22.57
10	78	6.43	29.00
11	68	5.60	34.60
12	72	5.93	40.53
13	66	5.44	45.96
14	62	5.11	51.07
15	67	5.52	56.59
16	58	4.78	61.37
17	61	5.02	66.39
18	86	7.08	73.48
19	56	4.61	78.09
20	72	5.93	84.02
21	59	4.86	88.88
22	74	6.10	94.98
23	61	5.02	100.00
Total	1,214	100.00	

Categoria a parte, no necesaria?
 xq se incluye en "4+ food groups"

Table 11.6 Infant and young child feeding (IYCF) practices

Percentage of youngest children age 6-23 months living with their mother who are fed according to three IYCF feeding practices based on breastfeeding status, number of food groups, and times they are fed during the day or night preceding the survey, by background characteristics, Rwanda 2014-15

Background characteristic	Among breastfed children 6-23 months, percentage fed:				Among non-breastfed children 6-23 months, percentage fed:				Among all children 6-23 months, percentage fed:					
	4+ food groups ¹	Minimum meal frequency ²	Both 4+ food groups and minimum meal frequency	Number of breastfed children 6-23 months	Milk or milk products ³	4+ food groups ¹	Minimum meal frequency ⁴	With 3 IYCF practices ⁵	Number of non-breastfed children 6-23 months	Breast milk, milk, or milk products ⁶	4+ food groups ¹	Minimum meal frequency ⁷	With 3 IYCF practices	Number of all children 6-23 months
Age in months														
6-8	14.1	41.7	11.8	467	*	*	*	*	7	99.1	13.9	41.9	11.6	474
9-11	30.0	37.8	15.9	420	*	*	*	*	5	98.9	30.1	37.5	15.7	425
12-17	32.3	49.7	20.4	748	(47.2)	(51.8)	(51.7)	(14.7)	46	96.9	33.5	49.8	20.0	793
18-23	34.0	55.0	23.5	627	29.2	58.5	42.6	8.5	88	91.2	37.0	53.5	21.6	716
Sex														
Male	27.3	49.8	18.2	1,122	38.3	50.5	51.9	12.3	75	96.2	28.7	50.0	17.9	1,196
Female	29.8	44.8	19.0	1,140	31.2	55.3	37.7	7.0	72	95.9	31.4	44.4	18.3	1,212
Residence														
Urban	45.6	53.3	31.7	355	34.4	63.6	46.9	14.8	45	92.6	47.7	52.5	29.8	400
Rural	25.4	46.2	16.2	1,907	35.0	48.1	44.0	7.4	102	96.7	26.6	46.1	15.7	2,009
Province														
City of Kigali	45.7	54.4	32.0	246	(34.2)	(66.2)	(42.6)	(13.1)	35	91.9	48.3	53.0	29.7	281
South	27.1	48.2	18.0	510	(53.8)	(46.9)	(65.2)	(16.6)	22	98.1	27.9	48.9	17.9	532
West	21.0	39.0	11.1	559	(15.9)	(32.2)	(20.3)	(8.7)	24	96.5	21.4	38.2	11.0	583
North	32.6	53.9	22.7	323	*	*	*	*	20	96.3	34.0	53.5	21.8	344
East	27.7	47.9	18.4	624	(34.7)	(55.6)	(48.4)	(5.1)	45	95.6	29.6	47.9	17.5	669
Mother's education														
No education	18.6	36.3	10.1	328	*	*	*	*	12	97.5	19.4	36.5	9.8	340
Primary	26.9	47.1	17.0	1,630	31.2	46.9	41.9	9.2	101	96.0	28.1	46.8	16.5	1,731
Secondary and higher	48.4	60.1	36.5	304	(47.0)	(75.4)	(55.8)	(15.1)	33	94.9	51.0	59.7	34.4	337
Wealth quintile														
Lowest	15.4	37.0	7.8	575	(40.4)	(33.0)	(48.2)	(4.0)	26	97.5	16.1	37.5	7.6	601
Second	21.6	45.7	13.9	485	*	*	*	*	20	97.1	22.6	45.4	13.7	505
Middle	25.2	45.3	16.5	440	(19.4)	(48.7)	(37.3)	(3.4)	29	95.1	26.6	44.8	15.7	469
Fourth	41.7	56.4	28.6	400	*	*	*	*	20	96.6	42.2	55.3	28.1	420
Highest	48.5	58.3	33.7	362	45.4	67.1	53.7	12.7	53	93.1	50.9	57.7	31.0	414
Total	28.6	47.3	18.6	2,262	34.8	52.9	44.9	9.7	147	96.0	30.1	47.2	18.1	2,409

Note: Figures in parentheses are based on 25-49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

¹ Food groups: a. infant formula, milk other than breast milk, cheese or yogurt or other milk products; b. foods made from grains, roots, and tubers, including porridge and fortified baby food from grains; c. vitamin A-rich fruits and vegetables and red palm oil; d. other fruits and vegetables; e. eggs; f. meat, poultry, fish, shellfish and organ meats; g. legumes and nuts.

² For breastfed children, minimum meal frequency is receiving solid or semisolid food at least twice a day for infants age 6-8 months and at least three times a day for children age 9-23 months.

³ Includes two or more feedings of commercial infant formula, fresh, tinned, and powdered animal milk, and yogurt

⁴ For nonbreastfed children age 6-23 months, minimum meal frequency is receiving solid or semisolid food or milk feeds at least four times a day.

⁵ Nonbreastfed children age 6-23 months are considered to be fed with a minimum standard of three infant and young child feeding practices if they receive other milk or milk products at least twice a day, receive the minimum meal frequency, and receive solid or semisolid foods from at least four food groups not including the milk or milk products food group.

⁶ Breastfeeding, or not breastfeeding and receiving two or more feedings of commercial infant formula, fresh, tinned, and powdered animal milk, and yogurt

⁷ Children are fed the minimum recommended number of times per day according to their age and breastfeeding status as described in footnotes 2 and 4.

Table 11.6 Infant and young child feeding (IYCF) practices

Percentage of youngest children age 6-23 months living with their mother who are fed according to three IYCF feeding practices based on breastfeeding status, number of food groups, and times they are fed during the day or night preceding the survey, by background characteristics, Rwanda 2010

Background characteristic	Among breastfed children 6-23 months, percentage fed:				Among nonbreastfed children 6-23 months, percentage fed:				Among all children 6-23 months, percentage fed:					
	4+ food groups ¹	Minimum meal frequency ²	Both 4+ food groups and minimum meal frequency ³	Number of breastfed children 6-23 months	Milk or milk products ³	4+ food groups ¹	Minimum meal frequency ⁴	With 3 IYCF practices ⁵	Number of non-breastfed children 6-23 months	Breast milk or milk products ⁶	4+ food groups	Minimum meal frequency ⁷	With all 3 IYCF practices	Number of all children 6-23 months
Age in month														
6-8	9.5	45.7	9.3	411	*	*	*	*	6	99.2	9.4	45.7	9.2	417
9-11	25.3	38.0	12.8	404	*	*	*	*	12	98.4	25.9	38.6	13.1	416
12-17	27.2	51.4	18.2	714	(33.8)	(47.9)	(52.0)	(15.2)	42	96.3	28.4	51.5	18.0	756
18-23	32.4	62.4	24.0	672	17.1	28.4	40.5	7.8	111	86.3	31.9	59.3	21.7	783
Sex														
Male	24.4	50.2	17.5	1,088	17.8	29.1	44.9	10.9	81	94.3	24.7	49.9	17.1	1,168
Female	25.9	52.2	17.1	1,112	30.1	37.5	44.7	9.9	91	94.7	26.8	51.7	16.6	1,204
Residence														
Urban	46.5	57.3	34.0	231	(36.2)	(53.7)	(58.5)	(17.3)	30	92.7	47.4	57.4	32.1	260
Rural	22.6	50.5	15.4	1,970	21.9	29.3	41.9	8.9	142	94.7	23.1	50.0	15.0	2,112
Province														
City of Kigali	45.8	56.6	33.5	182	(37.8)	(66.1)	(65.7)	(23.9)	32	90.7	48.8	58.0	32.1	213
South	32.0	55.7	22.2	545	(28.6)	(39.3)	(39.9)	(12.0)	25	95.9	32.3	55.0	21.8	570
West	14.6	41.9	9.5	605	(15.4)	(15.5)	(29.5)	(2.9)	40	94.8	14.6	41.1	9.1	645
North	26.8	55.4	18.1	322	*	*	*	*	26	94.0	26.8	54.7	17.4	348
East	22.1	52.9	15.3	546	(22.6)	(27.9)	(45.1)	(7.8)	49	93.6	22.6	52.3	14.7	595
Education														
No education	15.0	42.4	10.4	383	(28.1)	(27.9)	(46.2)	(2.8)	29	95.0	15.9	42.7	9.9	412
Primary	24.8	51.6	16.9	1,635	17.1	25.8	37.7	8.3	118	94.4	24.9	50.7	16.3	1,753
Secondary and higher	49.5	66.4	36.2	182	(53.9)	(76.0)	(76.0)	(28.7)	25	94.4	52.7	67.6	35.3	207
Wealth quintile														
Lowest	17.1	46.8	11.5	572	*	*	*	*	22	96.5	17.1	45.7	11.3	594
Second	20.6	47.9	12.4	511	(17.0)	(20.7)	(35.2)	(8.2)	27	95.9	20.6	47.3	12.2	537
Middle	23.3	48.1	14.2	393	(14.6)	(27.0)	(39.6)	(11.7)	37	92.6	23.6	46.5	14.0	430
Fourth	24.9	57.4	18.7	403	(19.6)	(25.1)	(45.8)	(3.8)	35	93.6	24.9	56.5	17.5	437
Highest	49.1	60.6	37.6	322	46.5	58.6	71.5	17.0	51	92.6	50.5	62.1	34.8	374
Total	25.1	51.2	17.3	2,200	24.4	33.6	44.8	10.4	172	94.5	25.8	50.8	16.8	2,372

Note: Figures in parentheses are based on 25-49 unweighted cases. An asterisk indicates that a figure is based on fewer than 25 unweighted cases and has been suppressed.

¹ Food groups: a. infant formula, milk other than breast milk, cheese or yogurt or other milk products; b. foods made from grains, roots, and tubers, including porridge and fortified baby food from grains; c. vitamin A-rich fruits and vegetables (and red palm oil); d. other fruits and vegetables; e. eggs; f. meat, poultry, fish, and shellfish (and organ meats); g. legumes and nuts.

² For breastfed children, minimum meal frequency is receiving solid or semisolid food at least twice a day for infants 6-8 months and at least three times a day for children 9-23 months.

³ Includes two or more feedings of commercial infant formula; fresh, tinned, and powdered animal milk; and yogurt.

⁴ For nonbreastfed children age 6-23 months, minimum meal frequency is receiving solid or semisolid food or milk feeds at least four times a day.

⁵ Nonbreastfed children age 6-23 months are considered to be fed with a minimum standard of three infant and young child feeding practices if they receive other milk or milk products at least twice a day, receive the minimum meal frequency, and receive solid or semisolid foods from at least four food groups not including the milk/milk product group.

⁶ Breastfeeding, or not breastfeeding and receiving two or more feedings of commercial infant formula; fresh, tinned, and powdered animal milk; and yogurt.

⁷ Children are fed the minimum recommended number of times per day according to their age and breastfeeding status as described in notes 2 and 4.