
Subject: Timeliness of measles vaccination

Posted by [nkanagat](#) on Mon, 11 Jul 2016 14:26:01 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hello,

I am trying to calculate the timeliness of the measles vaccination in Tanzania from the DHS 2010 dataset, KR file. I need to calculate whether the measles vaccine was given early (before 9 months) on time (9 months) or late (after 9 months).

I have set up the following code:

```
**Healthcard
```

```
gen healthcard=.
```

```
replace healthcard = 1 if h1 == 1
```

```
replace healthcard = 0 if inlist(h1, 0, 2, 3)
```

```
replace healthcard = . if h1 == 9
```

```
*generate age of each child in months
```

```
g age=v008-b3
```

```
g wgt=v005/1000000
```

```
ta age if age>=12 & age<=23 & b5==1 [iw=wgt]
```

```
**Measles
```

```
recode h9 (0 2 3 8 9=0 "no") (1=1 "yes"), g(measles)
```

```
ta measles if age>=12 & age<=23 & b5==1 [iw=wgt]
```

```
**Timeliness by month
```

```
recode h9m ( 1 2 3 4 5 6 7=0 "early") (9=1 "timely") (10 11 12=2 "late"), g (timely)
```

```
ta measles timely
```

In order to examine the timeliness of measles, is this the best way to set up my variables in stata?

Thank you,

Natasha