

---

Subject: Calculating malaria prevalence (Tanzania: Standard AIS, 2011-12)

Posted by [NinaB](#) on Mon, 21 Nov 2016 11:19:54 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hi,

I'm trying to reproduce the malaria prevalences (by mRDT and microscopy) for children 6-59 months which can be found in the Tanzania AIS 11/12 final report on page 157. I watched the DHS youtube videos on sampling and weighting and read some other threads in this forum. However, I was not able to get exactly the same numbers as in the report. I'm using the household member dataset (PR) and my stata code so far looks like this:

\*Parasitemia (via microscopy) in children 6-59 months

```
gen micmalpos=0
```

```
replace micmalpos=(hml32==1)
```

```
lab var micmalpos "Parasitemia (via microscopy) in children 6-59 months"
```

\*Parasitemia (via mRDT) in children 6-59 months

```
gen rdtmalpos=0
```

```
replace rdtmalpos=(hml35==1)
```

```
lab var rdtmalpos "Parasitemia (via mRDT) in children 6-59 months"
```

\*Creating weight variable

```
gen wgt=hv005/1000000
```

\*Survey setting the data

```
svyset [pw=wgt], psu(hv021) strata(hv023)
```

\*Proportion of children 6-59 months old with malaria infection (microscopy) by Region

```
svy: mean micmalpos if hc1>=6 & hc1<=59 & hml33==0, over(hv024)
```

\*Proportion of children 6-59 months old with malaria infection (mRDT) by Region

```
svy: mean rdtmalpos if hc1>=6 & hc1<=59 & hml33==0, over(hv024)
```

Does anyone have an idea what I'm doing wrong. Thanks for helping me.

Nina

---