
Subject: Re: Matching the sample size
Posted by Santosh on Wed, 18 Jul 2018 20:48:31 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Sammy,

I am sorry for late reply. Following are the codes I used.

```
// Systolic and Diastolic BP
gen sys = .
gen dia = .
* if the first measure is given
replace sys = shb16s if inrange(shb16s,20,299) & inrange(shb16d,20,299)
replace dia = shb16d if inrange(shb16s,20,299) & inrange(shb16d,20,299)
* if the second measure is given take it
replace sys = shb23s if inrange(shb23s,20,299) & inrange(shb23d,20,299)
replace dia = shb23d if inrange(shb23s,20,299) & inrange(shb23d,20,299)
* if the third measure is given take the last
replace sys = shb27s if inrange(shb27s,20,299) & inrange(shb27d,20,299)
replace dia = shb27d if inrange(shb27s,20,299) & inrange(shb27d,20,299)
* if all measures are given, then average the last two
replace sys = (shb23s + shb27s)/2 if inrange(shb27s,20,299) & inrange(shb27d,20,299) &
inrange(shb23s,20,299) & inrange(shb23d,20,299)
replace dia = (shb23d + shb27d)/2 if inrange(shb27s,20,299) & inrange(shb27d,20,299) &
inrange(shb23s,20,299) & inrange(shb23d,20,299)
```

Regards,
Sam
