

---

Subject: NFHS-4 Median Birth interval mismatch  
Posted by [dfnbcc](#) on Tue, 13 Aug 2019 17:03:46 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hello,

I am trying to calculate the median birth interval using STATA from the NFHS-4. However, I am getting the wrong median compared to the report. The report says the median in India is 32 months, however, I am getting 28 months.

I am subtracting 9 months from the variable as I want birth-to-pregnancy interval, not birth-to-birth. But even if I don't subtract 9, I get different values.

```
use IABR74FL.dta
gen weight=v005/1000000
```

```
rename sdistri shdistri
tostring shdistri, replace
gen strata = shdistri
svyset v021 [pweight=weight], strata (strata)
```

```
keep shdistri v008 b3 bord b0 b11 v106 hv024 v025 v190 v447a v133 s301 hv001 hv002 hvidx
v005 v021 weight strata
```

\*Births in the month of interview are included and births 60 months before the interview are excluded.

```
gen monthssncbirth=v008-b3
replace monthssncbirth=. if monthssncbirth<=60
drop if monthssncbirth==.
```

\*First-order births (and their twins) are excluded from both numerators and denominators.

```
svy: tab bord
gen birth_order=bord if (b0 > 1)
svy: tab birth_order
```

\*subtract by 9 months to find birth-to-pregnancy interval (WHO)

```
gen birth_interval=b11-9
replace birth_interval=. if birth_interval<0
```

---

Members Search Help Register Login Home Home » Topics » Fertility » NFHS-4  
Median Birth interval (Median birth interval does not match NFHS-4 report)  
Show: Today's Messages :: Show Polls :: Message Navigator  
Switch to threaded view of this topic Create a new topic Submit Reply  
- NFHS-4 Median Birth interval [message #17992] Sun, 11 August 2019 11:28  
dfnbcc is currently offline dfnbcc

Messages: 2  
Registered: June 2019  
Member  
Hello,

I am trying to calculate the median birth interval using STATA from the NFHS-4. However, I am getting the wrong median compared to the report. The report says the median in India is 32 months, however, I am getting 28 months.

I am subtracting 9 months from the variable as I want birth-to-pregnancy interval, not birth-to-birth. But even if I don't subtract 9, I get different values.

```
Code: [Select all] [Show/ hide]
use IABR74FL.dta
gen weight=v005/1000000
```

```
rename sdistri shdistri
tostring shdistri, replace
gen strata = shdistri
svyset v021 [pweight=weight], strata (strata)
```

```
keep shdistri v008 b3 bord b0 b11 v106 hv024 v025 v190 v447a v133 s301 hv001 hv002 hvidx
v005 v021 weight strata
```

\*Births in the month of interview are included and births 60 months before the interview are excluded.

```
gen monthssncbirth=v008-b3
replace monthssncbirth=. if monthssncbirth<=60
drop if monthssncbirth==.
```

\*First-order births (and their twins) are excluded from both numerators and denominators.

```
svy: tab bord
gen birth_order=bord if (b0 > 1)
svy: tab birth_order
```

\*subtract by 9 months to find birth-to-pregnancy interval (WHO)

```
gen birth_interval=b11-9
replace birth_interval=. if birth_interval<0
```

Would appreciate some help please.

Thanks in advance.