
Subject: What Should be Used "And (&)" "or "OR (|)"
Posted by shujaat.smc@gmail.com on Tue, 27 Oct 2020 09:08:52 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear DHS Representative,

Before performing descriptives and inferential analysis, after declaring my data set (BR file PDHS 2017-18). I run the following command for my univariate analysis;

```
svy linearized, subpop(if bidx==1 & b0==0 & v210!=1) : stcox ib(0). HepBTest
```

Quest: I am using subpopulation for those neonates who were last born to the mothers included in the survey (bidx==1), who was the product of singleton pregnancy (b0==0), and who were not born in the month of conducting the survey (v210!=1). Kindly confirm and please also explain is this correct to use "AND (&)" in between my subpopulation variables. This query pops up in my mind after seeing the coding on the git-hub "

https://github.com/DHSProgram/DHS-Indicators-Stata/blob/master/Chap09_RH/RH_DEL.do
over here the "OR (|)" is used.

Quest: What is meant by "if age>=period" I am asking this question because in my analysis up till now I have confidence in my sup-population, however after seeing the above git-hub link I in doubt regarding AND(&) OR(|) use. Also, I have not used any stata command if age>=period up till now.

I am working on the determinants of neonatal mortality in Pakistan (BR-file 2017-18) using STATA 13.1.

Waiting for your reply.

Best Regards

Dr. Hussain