Subject: Re: Continuous Survey Weights

Posted by Bridgette-DHS on Wed, 04 Jun 2014 13:57:44 GMT

View Forum Message <> Reply to Message

Following is a response from Senior DHS Specialist, Shea Rutstein:

The Peru Continuous DHS started its first cycle in 2004. From 2004 to 2007, the yearly sample was a 1/5th random sample of the clusters drawn for the 2000 Peru DHS, which was based 1999 pre-census sampling frame (the census never was realized. This sub-selection was done to reduce sampling errors for trends. Thus the principal component of the sample weights were the selection probabilities for those clusters in 2000. Adjustments were made to the weights for three items: the change in the number of dwellings between 2000 and the cycle year (a relisting of each cluster was performed before interviewing fieldwork); non-response rates for households and women; and post-stratification for better representation (given that it was a random subsample of a sample). In 2008, an additional large set of clusters was added to increase sample size. A different frame was used for the additional clusters and weights were calculated for the combined frames to represent urban and rural areas of each department (state) according to the census of 2007, as well as adjusting for non-response. Given that none of the clusters was reused, the sample weights can be used for individual years. From 2009 onward, the sample frame is based on the 2007 census, so that the sampling weights can also be used for comparisons without problems, even though 1/2 of the clusters of each year's sample was reused in the following year to reduce the sampling errors for trends.

Anthropometric measurements were taken in 2005, 2007, and from 2008 onward. From 2008, the survey recorded the height and weight of children under five years that was noted on their vaccination cards during Growth and Development Controls. These data include the date of the control and can serve as real cohort information. You may want to use them instead.