

---

Subject: Re: Penta1 to measles dropout

Posted by [Janet-DHS](#) on Tue, 10 May 2022 14:23:06 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Janet-DHS wrote on Tue, 10 May 2022 10:22: Following is response from DHS Research & Data Analysis Director, Tom Pullum:

All the numbers in DHS reports are weighted by v005. Are you using weights? If you cannot match a number in a report, please tell us which number in which table in which report.

I have not worked with a DPT dropout variable before but I assume your DPT\_dropout variable should be calculated as follows:

```
gen DPT_dropout=0 if inrange(h3,1,3)==1  
replace DPT_dropout=1 if DPT_dropout==0 & inrange(h,1,3)==0
```

I don't know what your "ms" variable is (it must be related to h9) but the logic would be the same. The dropout variable should be a binary variable that is conditional on having received DPT1, and it is NA (a dot in Stata) if the child did not receive DPT1. That is, it is NA if the child did not receive DPT1, 0 if it received DPT1 but not DPT3, and 1 if it received DPT1 AND DPT3. That's what the two Stata lines would construct.

The woman is interviewed. In the KR file there is one record for each child born in the past five years. The file includes children who have died. The child's record includes most of the mother's variables, and there is hardly ever a reason to merge the KR and IR files.