
Subject: Re: Identifying siblings of a child that are below 1 years of age (R)

Posted by [Janet-DHS](#) on Mon, 01 Aug 2022 16:25:32 GMT

[View Forum Message](#) <> [Reply to Message](#)

Following is a response from DHS Research & Data Analysis Director, Tom Pullum:

Usually we at DHS would not answer a question like this, but I have always been interested in kinship and decided to work on it.

You did not say what survey you are using. I will use the Bangladesh 2017-18 survey (BD7R) to illustrate.

Two children are siblings if they have the same mother. I will re-state your question to be as follows: "For a child age A or less, how many YOUNGER siblings do they have who are age B or less." You described this with B=0 but I am generalizing. In my example, A is 17 and B is 0, 1, 2, 3, or 4. I assume the question is about living children and siblings but that could be changed.

Usually two children the same age will be twins, but I allow for that. You have to be sure not to count a child as its own sibling. The ages of living children are b8. You have to store a temporary file.

Open the BR file (BDBR7RFL.dta) in Stata. Enter the lines pasted below. Good luck.

```
cd e:\DHS\DHS_data\scratch
```

```
use "C:\Users\26216\ICF\Analysis - Shared Resources\Data\DHSdata\BDBR7RFL.DTA" , clear  
keep if b5==1
```

```
keep v001 v002 v003 bidx b8
```

```
forvalues li=0/4{  
gen nsibs_age_le_`li'=0  
}
```

```
sort v001 v002 v003
```

```
save index_children.dta, replace
```

```
drop nsibs*  
rename bidx bidx_sib  
rename b8 b8_sib
```

```
sort v001 v002 v003  
joinby v001 v002 v003 using index_children.dta
```

```
forvalues li=0/4{  
replace nsibs_age_le_`li'=nsibs_age_le_`li'+1 if bidx_sib<bidx & b8_sib<=`li'  
}
```

```
drop bidx_sib b8_sib
```

```
collapse (sum) nsibs_age_le_* (first) b8, by(v001 v002 v003 bidx)
```

```
forvalues li=0/4{
```

```
tab b8 nsibs_age_le_`li' if b8<=17
```

```
}
```
