
Subject: Re: anaemia in children

Posted by [Sadhvi](#) on Wed, 30 Nov 2022 06:00:04 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi.

The following is the code in r that I am using.

```
hh_recode = read_sas("Path\\IAPR7BFL.SAS7BDAT")
```

```
# Here I am filtering out the children who slept in the household last night, and aged 6 to 59 months and have a non-NA value for the anaemia status - HC57
```

```
cleanedAneSubsetNfhs5 = hh_recode[hh_recode$HV103 == 1 & !is.na(hh_recode$HC1) & hh_recode$HC1 > 5 & hh_recode$HC1 < 60 & !is.na(hh_recode$HC57),]
```

```
#overall anaemia
```

```
weightedSample = sum(cleanedAneSubsetNfhs5$HV005)/ 1000000
```

```
#now filtering out the children who have mild, moderate or severe anaemia as given by the variable HC57.
```

```
WeightedAnChildren = sum(cleanedAneSubsetNfhs5$HV005[cleanedAneSubsetNfhs5$HC57 %in% c(1,2,3)]) / 1000000
```

```
anemiaPrevalence = WeightedAnChildren/weightedSample
```

```
# Results: The total sample that turns out is 178952.3 but the number given in table 10.12 of the national report is 152,752. And the prevalence of anemia in children 6 to 59 months is 68.07%
```
