

---

Subject: Re: Replicating Step 7 and Level-2 Weight from Report 27

Posted by [Yawo](#) on Wed, 12 Jul 2023 21:20:55 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Thanks for your response. I understand the procedure, so just to be clear, I have two related questions:

Q1: Given that I am combining the data for both men and women, are the following codes ok now?

```
***** STAGE A *****
```

```
* 7: Prepare the adjusted final/individual adjusted weight for men and women: dividing population of women or men by the appropriate number interviewed; de-normalize the final weight */
```

```
gen newwt =.  
replace newwt =(v005/1000000)*(4764000/10060) if sex==1  
replace newwt =(v005/1000000)*(5538000/5135) if sex==2  
label variable newwt "Population adjusted sample weight"
```

```
***** STAGE B: APPROXIMATE LEVEL-1 AND LEVEL-2 WEIGHTS *****
```

```
/*Step 1: De-normalize the final weight, using approximated normalization factor. */
```

```
gen d_HH = newwt * (M/m_c)
```

```
/* Step 2: Approximate the level-2 weight (get f variation factor; set alpha based equal split
```

```
gen f = d_HH / ((A_h/a_c_h) * (M_h/S_h))  
scalar alpha = 0.5  
gen wt2 = (A_h/a_c_h)*(f^alpha)
```

```
/* Step 2B - OPTIONAL: Generate a revised level-2 weight by replacing it with its geometric mean [if needed to get resolve the "weights in variable wt2 not constant within groups defined by: v001" error - based on advice by Tom here in [message
```

```
#21803]:https://userforum.dhsprogram.com/index.php?t=msg& ;amp ;amp ;amp ;th=9098&goto=21803&#msg\_21803
```

```
gen ln_wt2=log(wt2)  
egen mean_wt2=mean(ln_wt2), by(v001)  
gen rev_wt2=exp(mean_wt2)  
drop ln_wt2
```

```
/* Step 3: Approximate the level-1 weight */ gen wt1 = d_HH/wt2
```

Q2: If rev\_wt2 is used, then should we adjust the formula for wt1 as well: gen rev\_wt1 = d\_HH/rev\_wt? By the same token, should we also change to the rev\_wt2 in the loop (i.e, gen

wt1\_`i' = d\_HH/wt2\_`i')?

With much appreciation for your assistance.

Cy

---