
Subject: KDHS 2014: Table 2.14 School attendance ratios

Posted by [sokiya](#) on Tue, 07 Nov 2023 17:11:46 GMT

[View Forum Message](#) <> [Reply to Message](#)

I am trying to generate Table 2.14 School attendance ratios using the microdata and the code from the DHS GitHub repo as shown below

* open the birth history data to extract date of birth variables needed.

use "KEBR71FL.DTA", clear

* keep only the variables we need

keep v001 v002 v003 b3 b16

* drop if the child in the birth history was not in the household or not alive

drop if b16==0 | b16==.

* rename key variables for matching

rename b16 hvidx

rename v001 hv001

rename v002 hv002

* sort on key variables

sort hv001 hv002 hvidx

* if there are some duplicates of line number in household questionnaire, we need to drop the duplicates

gen dup = (hv001 == hv001[_n-1] & hv002 == hv002[_n-1] & hvidx == hvidx[_n-1])

drop if dup==1

drop dup

* re-sort to make sure still sorted

sort hv001 hv002 hvidx

* save a temporary file for merging

tempfile tempBR

save `tempBR'

* use the PR file for household members for the NAR and GAR indicators

use "KEPR71FL.DTA", clear

* merge in the date of birth from the women's birth history for the household member

merge 1:1 hv001 hv002 hvidx using `tempBR'

* there are a few mismatches of line numbers (typically a small number of cases) coming from the BR file, so let's drop those

drop if _merge==2

* restrict to de facto household members age 5-24, and drop all others

keep if hv103==1 & inrange(hv105,5,24)

* now we calculate the child's age at the start of the school year

* but first we have to specify the month and year of the start of the school year referred to in the

survey

* example, for Zimbabwe 2015 survey this was January 2015

global school_start_yr = 2014

global school_start_mo = 1

* also need the age ranges for primary and secondary

global age_prim_min = 6

global age_prim_max = 13

global age_sec_min = 14

global age_sec_max = 17

* produce century month code of start of school year for each state and phase

gen cmcSch = (\$school_start_yr - 1900)*12 + \$school_start_mo

replace cmcSch = cmcSch+12 if hv008 >= cmcSch+12

* calculate the age at the start of the school year, using the date of birth from the birth history if we have it

gen school_age = int((cmcSch - b3) / 12) if b3 != .

* Impute an age at the beginning of the school year when CMC of birth is unknown

* the random imputation below means that we won't get a perfect match with the report, but it will be close

gen xtemp = hv008 - (hv105 * 12) if b3 == .

gen cmctemp = xtemp - int(uniform()*12) if b3 == .

replace school_age = int((cmcSch - cmctemp) / 12) if b3 == .

* Generate variables for whether the child is in the age group for primary or secondary school

gen prim_age = inrange(school_age,\$age_prim_min,\$age_prim_max)

gen sec_age = inrange(school_age,\$age_sec_min,\$age_sec_max)

* create the school attendance variables, not restricted by age

gen prim = (hv122 == 1)

gen sec = (hv122 == 2)

* set sample weight

cap gen wt = hv005/1000000

* For NAR we can use this as just regular variables and can tabulate as follows, but can't do this for GAR as the numerator is not a subset of the denominator

* NAR is just the proportion attending primary/secondary school of children in the correct age range, for de facto children

gen nar_prim = prim if prim_age == 1

gen nar_sec = sec if sec_age == 1

lab var nar_prim "Primary school net attendance ratio (NAR)"

lab var nar_sec "Secondary school net attendance ratio (NAR)"

* tabulate primary school attendance

tab hv104 nar_prim [iw=wt] , row

tab hv025 nar_prim [iw=wt] , row

tab hv270 nar_prim [iw=wt] , row

* tabulate secondary school attendance

```
tab hv104 nar_sec [iw=wt] , row
tab hv025 nar_sec [iw=wt] , row
tab hv270 nar_sec [iw=wt] , row
```

Any help will be appreciated
