

---

Subject: Help-Seeking Behaviour of Domestic Violence\_Myanmar DHS (2015-16)

Posted by [Soe Myat Htet](#) on Wed, 31 Jul 2024 11:26:48 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Dear Moderator,

May I discuss two questions regarding "Weighting"?

(1) I am doing DHS data analysis using "R". Before going data analysis, I did weight the selected variables after managing missing values and mutating some variables. Then, I compared the results of unweighted and weighted frequencies. The unusual differences in 7 categories were found in one variable "v101" which had 15 categories. For example, (n = 60) before weight and (n = 5.11) after weight. I checked again and again. Also, I checked in Stata. It showed similar. Is it possible? Should I weight or not? I used R codes as below:

```
IRdata$wt<- IRdata$v005/1000000
mysurvey <- svydesign(id = IRdata$v021, data = IRdata, strata = IRdata$v022,
                    weights = IRdata$wt, nest = T)
options (survey.lonely.psu = "adjust")
svytable(~ v101, mysurvey)
```

(2) In relation to weighting logistic regression analysis. Some researchers say that it should be weighted all analysis (descriptive, bivariate, and multivariate analysis). However, some researchers say that it should be considered for descriptive data analysis only. Please kindly give me your suggestion.

Thank you,  
Soe