Subject: Re: variables for svyset in Stata (Bangladesh 2011) Posted by mmr-UMICH on Wed, 04 Feb 2015 03:58:29 GMT View Forum Message <> Reply to Message

## Thank you, Trevor.

I will cut and paste some lines whenever requires from original message and try to clarify concretely the reasons behind that error/warnings and its possible solution:

\*\*now analyzing using poisson for the lower 2 wealth quintiles\*\* svy: poisson csect v190 if v190<3

The above svy: command is not recommended [and not correctly handle the domain concept] as "if v190<3" in svy: subsets the data (i.e., also deleting design information [aside: full sample design information is important for correctly calculating the sampling errors]) prior to run the poisson regression.

This svy: command uses such a subset data and as a result analysis sample (i.e. "estimation sample" in Stata wording) lacks one strata and 101 PSUs (see below output (cut and paste): -----start----

```
(running poisson on estimation sample)
```

Survey: Poisson regression

Number of strata = 19 Number of obs = 3645 Number of PSUs = 499 Population size = 3867.0893 Design df = 480 ------ end -----

We have to create a variable, say: mydomain = 1 if v190 < 3, otherwise, mydomain = 0, then use svy command: svy, subpop(mydomain): poisson csect v190

I hope this run will not encounter such issue and also does not require singleunit(centered)\* svyset option. And the output will show the same # of obs and population size, but changed others stats such as # of strata, PSUs and degrees of freedom (df).

I verified that strata 5 and 11 have 5 and 23 PSUs respectively; so this svy, subpop(): that form "analytic" domain/subpopulation/subgroup will not be an issue of singleton-strata from full sample data.

\*note that singleunit(method) is kind of practically recommended for "analytic" subgroup and/or subclass analysis which sometimes encounter singleton-strata. This specification also appropriately calculates the degrees of freedom, which is prim important for statistical inferences, e.g, confidence intervals and p-values estimation.

Thank you all again.

Moshiur Rahman