
Subject: Re: Comparative Wealth Index for Bangladesh
Posted by [mmr-UMICH](#) on Sun, 21 Feb 2016 05:44:42 GMT
[View Forum Message](#) <> [Reply to Message](#)

Thank you so much, Dr. Trevor.

I have a related question for my previous request. You know I wanted to calculate the rural wealth index for Bangladesh DHS 2011, as I mentioned previous email that I downloaded an SPSS syntax file from DHS website which QH series variables using DHS raw data. I am trying to prepare the code as per your advice; trying to find the corresponding variables of QH series from HV variables in recode data set. I did almost major part of the syntax but I cannot figure out corresponding HV variables for the following variables (I cut and paste from the downloaded syntax code). Could you please provide me the variable labels of these variables at your earliest possible convenience. Thanking you again.

(I collect the variables here: qh110A to qh110L qh118A to qh118D qh121 qh122a to qh122c qh122_1 to qh122_5 qh123 and DOMESTIC)

```
*{Reset missing values to "does not have", change 2 code to 0}.
```

```
if (missing(qh110a) | qh110a<>1) qh110a=0.  
if (missing(qh110b) | qh110b<>1) qh110b=0.  
if (missing(qh110c) | qh110c<>1) qh110c=0.  
if (missing(qh110d) | qh110d<>1) qh110d=0.  
if (missing(qh110e) | qh110e<>1) qh110e=0.  
if (missing(qh110f) | qh110f<>1) qh110f=0.  
if (missing(qh110g) | qh110g<>1) qh110g=0.  
if (missing(qh110h) | qh110h<>1) qh110h=0.  
if (missing(qh110i) | qh110i<>1) qh110i=0.  
if (missing(qh110j) | qh110j<>1) qh110j=0.  
if (missing(qh110k) | qh110k<>1) qh110k=0.  
if (missing(qh110l) | qh110l<>1) qh110l=0.  
if (missing(qh118a) | qh118a<>1) qh118a=0.  
if (missing(qh118b) | qh118b<>1) qh118b=0.  
if (missing(qh118c) | qh118c<>1) qh118c=0.  
if (missing(qh118d) | qh118d<>1) qh118d=0.  
if (not(missing(qh122c)) & qh122c < 99.98) landarea=qh122c.  
if (qh122c=99.95) landarea=95.  
if (missing(qh122b) | qh122b<>1) landarea=0.  
frequencies landarea.  
if (missing(qh121) | qh121 <>1) qh121=0.  
if (missing(qh122_1) | qh121 <>1) qh122_1=0.  
if (missing(qh122_2) | qh121 <>1) qh122_2=0.  
*if (missing(qh122_3) | qh121 <>1) qh122_3=0.  
if (missing(qh122_4) | qh121 <>1) qh122_4=0.  
if (missing(qh122_5) | qh121 <>1) qh122_5=0.  
if (missing(qh122a) | qh122a<>1) qh122a=0.  
missing values qh122_1 to qh122_5 (98,99).  
if (missing(qh123) | qh123<>1) qh123=0.
```

