

---

Subject: Re: Data Analysis in SPSS

Posted by [Trevor-DHS](#) on Fri, 06 Dec 2013 15:34:33 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

First, I moved your post from the "Dataset Use in Stata" forum to the "Dataset Use in SPSS" forum as it is about using data in SPSS.

In response to your questions:

1) To match the numbers in the DHS reports, first check that you are getting the same denominator as DHS. For women and men, the denominator is all de facto women and men who were tested for HIV - check for HIV03 in the range 0:7 for each man or women. The weight to use is HIV05/1000000. Once you are matching the denominator, then the numerator is a subset of these cases, with HIV03 = 1 or 3 (we only report those cases positive with HIV-1).

2) Complex Samples procedures are not required to match the estimates of HIV prevalence. The Complex Samples procedures are necessary if you are producing sampling errors, confidence intervals or testing for significance. Using Complex Sample does not affect the basic calculation of the indicator. If you do not use Complex Samples procedures, your software is assuming that you are using a simple random sample and all of your estimates of sampling errors, confidence intervals and your significance tests would be wrong. Complex Samples is an additional module that needs to be ordered for SPSS. If you have not ordered this module, then it will not appear in your list of available procedures in SPSS.

3) Once you set up a Complex Samples plan for your analysis, you then use this plan with the Complex Samples procedures such as CSDESCRIPTIVES, CSTABULATE, CSLOGISTIC, etc. to run your analysis. These procedures differ a little from the regular SPSS procedures that you may be familiar with, but generally you can produce the equivalent results with the Complex Samples procedures, but taking into account the sample design.

I hope this helps.