

---

Subject: Denormalization of weights required?

Posted by [MiFoo](#) on Sun, 02 May 2021 21:28:09 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hello,

I am using the BDHS from 2011 and 2017/2018 (PR file) to estimate a model of the form  $\text{svyglm}(\text{outcome} \sim \text{group} * \text{year} + \text{controls}, \text{design} = \text{sdata}, \text{family} = \text{binomial})$  followed by the calculation of average marginal effects of the change in the outcome over time for each group.

I ensured that that strata and PSU ID codes are unique in both surveys before creating the pooled survey design object by adding a year suffix. However, I am not sure whether a denormalization of weights is required for this kind of analysis. Some people in the User Forum said that different scales of weights in single surveys should not matter if I look at the results by year (i.e. include a year dummy). But multiplying the weights in 2011 by a random number changed my estimated regression coefficients (including the interaction term) and the marginal effects even though not very much?

Do I need to denormalize the weights and if yes is it right to multiply the weights by  $(\text{total number of residential households in the country in the year of the survey round}) / (\text{total number of households interviewed in the survey})$  regardless of the fact that 1) I am looking at individuals not households and 2) I am using only a subset of people (e.g. age>40)

Best,  
MiFoo

---

---

Subject: Re: Denormalization of weights required?

Posted by [Bridgette-DHS](#) on Mon, 03 May 2021 13:02:52 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Following is a response from DHS Senior Sampling Specialist, Mahmoud Elkasabi:

I would advise you to de-normalize the weight of the two survey. This would be the safest approach to use. It is fine to multiply the weight by  $(\text{total number of residential households in the country in the year of the survey round}) / (\text{total number of households interviewed in the survey})$ . It would be even better to multiply the weight by  $(\text{total number of adults 40+ in the country in the year of the survey round}) / (\text{total adults 40+ listed in the survey})$ .

---