

---

Subject: Post Natal Care check

Posted by [Francois](#) on Tue, 30 Nov 2021 07:34:35 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hello,

I am trying to reproduce the Table 9.9 and Table 9.11 ,The percentage of women with a pnc chek during the first 2days after birth for both mothers and newborn using GITHUB Codes. I am getting the different numbers in frequencies and percentages, Probably variables used changed but don't know where are mistakes. Thank you for your assistance

Here are the codes used:

```
*=====PNC INDICATORS=====
```

```
cd"C:\Users\ICF Rwanda\Desktop\district\DHS6"
```

```
use RWIR81FL, clear
```

```
gen age = v008 - b3_01
```

```
gen w=v005/1000000
```

```
scalar drop _all
```

```
** To check if survey has m51_1, which was in the surveys before 2005.
```

```
scalar m51_included=1
```

```
capture confirm numeric variable m51_1, exact
```

```
if _rc>0 {
```

```
  * m51_1 is not present
```

```
  scalar m51_included=0
```

```
}
```

```
if _rc==0 {
```

```
  * m51_1 is present; check for values
```

```
  summarize m51_1
```

```
    if r(sd)==0 | r(sd)==. {
```

```
      scalar m51_included=0
```

```
    }
```

```
}
```

```
*** Mother's PNC ***
```

```
if m51_included==1 {
```

```
//PNC timing for mother
```

```
recode m51_1 (100/103 = 1 "<4hr") (104/123 200 = 2 "4-23hrs") (124/171 201/202 = 3 "1-2 days") ///
```

```
(172/197 203/206 = 4 "3-6 days") (207/241 301/305=5 "7-41 days") ///
```

```
(198/199 298/299 398/399 998/999 = 9 "dont know/missing") (242/297 306/397 = 0 "no pnc check") , g(rh_pnc_wm_timing)
```

```
replace rh_pnc_wm_timing = 0 if m50_1==0 | m50_1==9
```

```
replace rh_pnc_wm_timing = 0 if (m52_1>29 & m52_1<97) | m52_1==.
```

```

replace rh_pnc_wm_timing=. if age>=24 | bidx_01!=1
label var rh_pnc_wm_timing "Timing after delivery for mother's PNC check"

```

```

//PNC within 2days for mother
recode rh_pnc_wm_timing (1/3= 1 "Visit w/in 2 days") (0 4 5 9 = 0 "No Visit w/in 2 days"),
g(rh_pnc_wm_2days)
label var rh_pnc_wm_2days "PNC check within two days for mother"

```

```

//PNC provider for mother
** This is country specific and could be different for different surveys, please check footnote of the
table for this indicator in the final report.
recode m52_1 (0 = 0 "No check") (11 = 1 "Doctor") (12/13 = 2 "Nurse/Midwife") ( 14/15 = 3 "Other
skilled provider") (16/90 = 4 "Non-skilled provider") (96 = 5 "Other") ///
( else = 9 "Don't know or missing") if age<24 & rh_pnc_wm_2days==1, gen(rh_pnc_wm_pv)
replace rh_pnc_wm_pv = 0 if rh_pnc_wm_2days==0 & age<24
label var rh_pnc_wm_pv "Provider for mother's PNC check"
}

```

```

if m51_included==0 {
cap drop rh_pnc_wm_timing

```

```

//PNC timing for mother
*did the mother have any check
gen momcheck = 0 if age<24
replace momcheck = 1 if (m62_1==1 | m66_1==1) & age<24

```

```

*create combined timing variable
gen pnc_wm_time = 999 if (age<24 & momcheck==1)
*start with women who delivered in a health facility with a check
replace pnc_wm_time = m63_1 if inrange(m64_1,11,29) & age<24
*Account for provider of PNC- country specific- see table footnotes
replace pnc_wm_time = 0 if (pnc_wm_time < 1000 & m64_1 >30 & m64_1 < 100 & age<24)
*Add in women who delivered at home with a check
replace pnc_wm_time = m67_1 if (pnc_wm_time == 999 & inrange(m68_1, 11,29) & age<24)
*Account for provider of PNC- country specific- see table footnotes
replace pnc_wm_time = 0 if m67_1 < 1000 & m68_1 >30 & m68_1 < 100 & age<24
*Add in women who had no check
replace pnc_wm_time = 0 if momcheck == 0 & age<24

```

```

*Recode variable into categories as in FR
recode pnc_wm_time (0 242/299 306/899 = 0 "No check or past 41 days") ( 100/103 = 1 "<4hrs")
(104/123 200 = 2 "4-23hrs") (124/171 201/202 = 3 "1-2days") ///
(172/197 203/206 = 4 "3-6days") (207/241 301/305 = 5 "7-41days") (else = 9 "Don't
know/missing") if age<24, gen(rh_pnc_wm_timing)
*label variable
label var rh_pnc_wm_timing "Timing after delivery for mother's PNC check"

```

```

//PNC within 2days for mother

```

```

recode rh_pnc_wm_timing (1/3 = 1 "Within 2 days") (0 4 5 9 = 0 "Not in 2 days"),
gen(rh_pnc_wm_2days)
label var rh_pnc_wm_2days "PNC check within two days for mother"

```

```
//PNC provider for mother
```

**\*\* This is country specific and could be different for different surveys, please check footnote of the table for this indicator in the final report.**

**\*Providers of PNC for facility deliveries**

```

recode m64_1 (0 = 0 "No check") (11 = 1 "Doctor") (12/13 = 2 "Nurse/Midwife") ( 14/15 = 3 "Other
skilled provider") (16/90 = 4 "Non-skilled provider") (96 = 5 "Other") ///
( else = 9 "Don't know or missing") if age<24 & rh_pnc_wm_2days==1, gen(pnc_wm_pv_hf)
replace pnc_wm_pv_hf = 0 if rh_pnc_wm_2days==0 & age<24

```

**\*Providers of PNC for home deliveries or checks after discharge**

```

recode m68_1 (0 = 0 "No check") (11 = 1 "Doctor") (12/13 = 2 "Nurse/Midwife") ( 14/15 = 3 "Other
skilled provider") (16/90 = 4 "Non-skilled provider") (96 = 5 "Other") ///
( else = 9 "Don't know or missing") if age<24 & rh_pnc_wm_2days==1 , gen(pnc_wm_pv_home)
replace pnc_wm_pv_home = 0 if rh_pnc_wm_2days==0 & age<24

```

**\*Combine two PNC provider variables**

```

clonevar rh_pnc_wm_pv = pnc_wm_pv_hf
replace rh_pnc_wm_pv = pnc_wm_pv_home if (pnc_wm_pv_hf==9 & rh_pnc_wm_2days==1 &
age<24)
*label variable
label var rh_pnc_wm_pv "Provider for mother's PNC check"
}

```

**\*\*\* Newborn's PNC \*\*\***

**\* some surveys (usally older surveys) do not have PNC indicators for newborns. For this you would need variables m70\_1, m71\_1, ..., m76\_1**

```

scalar m70s_included=1
forvalues i=1/6 {
  capture confirm numeric variable m7`i'_1, exact
  if _rc>0 {
    scalar m70s_included=0
  }
  if _rc==0 {
    summarize m7`i'_1
    if r(sd)==0 | r(sd)==. {
      scalar m70s_included=0
    }
  }
}
}

```

```

*survey has newborn PNC indicators
if m70s_included==1 {

  if m51_included==1 {

    //PNC timing for newborn
    recode m71_1 (207/297 301/397 = 0 "No check or past 7 days") ( 100 = 1 "less than 1 hour")
    (101/103 = 2 "1-3 hours") (104/123 200 = 3 "4 to 23 hours") (124/171 201/202 = 4 "1-2 days") ///
    (172/197 203/206 = 5 "3-6 days new") (198/199 298/299 398/399 998/999 = 9 "dont
    know/missing") if age<24 , gen(rh_pnc_nb_timing)

    *Recode babies with no check and babies with check by unskilled prov back to 0
    replace rh_pnc_nb_timing = 0 if (m70_1==0 | m70_1==9)
    *Account for provider of PNC- country specific- see table footnotes
    replace rh_pnc_nb_timing = 0 if (m72_1>29 & m72_1<97) | m72_1==.
    replace rh_pnc_nb_timing = . if age>=24 | bidx_01!=1

    *label variable
    label var rh_pnc_nb_timing "Timing after delivery for newborn's PNC check"

    //PNC within 2days for newborn
    recode rh_pnc_nb_timing (1/4 = 1 "Visit within 2 days") (0 5 9 = 0 "No Visit within 2 days"),
    g(rh_pnc_nb_2days)
    *label variable
    label var rh_pnc_nb_2days "PNC check within two days for newborn"

    //PNC provider for newborn
    ** this is country specific, please check table in final report
    recode m72_1 (0 = 0 "No check") (11 = 1 "Doctor") (12/13 = 2 "Nurse/Midwife") ( 14/15 = 3
    "Other skilled provider") (16/90 = 4 "Non-skilled provider") (96 = 5 "Other") ///
    ( 98/99 = 9 "Don't know or missing") if age<24 & rh_pnc_nb_timing<9 & rh_pnc_nb_timing>0,
    gen(rh_pnc_nb_pv)
    replace rh_pnc_nb_pv = 0 if rh_pnc_nb_2days ==0 & age<24
    label var rh_pnc_nb_pv "Provider for newborn's PNC check"

  }

  if m51_included==0 {

    //PNC timing for newborn

    *Newborn check
    gen nbcheck = 1 if (m70_1==1 | m74_1==1 )
    *create combined timing variable
    gen pnc_nb_timing_all = 999 if age<24 & nbcheck==1

    *start with women who delivered in a health facility with a check
    replace pnc_nb_timing_all = m75_1 if inrange(m76_1,11,29) & age<24
  }
}

```

\*Account for provider of PNC- country specific- see table footnotes  
replace pnc\_nb\_timing\_all = 0 if pnc\_nb\_timing\_all < 1000 & m76\_1 >30 & m76\_1 < 100 & age<24

\*Add in women who delivered at home with a check  
replace pnc\_nb\_timing\_all = m71\_1 if (pnc\_nb\_timing\_all==999 & inrange(m72\_1,11,29) & age<24)

\*Account for provider of PNC- country specific- see table footnotes

replace pnc\_nb\_timing\_all = 0 if (m71\_1 < 1000 & m72\_1 >30 & m72\_1 < 100 & age<24)

\*Add in women who had no check

replace pnc\_nb\_timing\_all = 0 if (nbcheck!=1) & age<24

\*Recode variable into categories as in FR

recode pnc\_nb\_timing\_all (0 207/297 301/397 = 0 "No check or past 7 days") ( 100=1 "less than 1 hour") (101/103 =2 "1-3 hours") (104/123 200 = 3 "4 to 23 hours") (124/171 201/202 = 4 "1-2 days") ///

(172/197 203/206 = 5 "3-6 days new") (else = 9 "Don't know or missing") if age<24 , gen(rh\_pnc\_nb\_timing)

\*label variable

label var rh\_pnc\_nb\_timing "Timing after delivery for mother's PNC check"

//PNC within 2days for newborn

recode rh\_pnc\_nb\_timing (1/4 = 1 "visit within 2 days") (0 5 9 = 0 "No Visit within 2 days"), g(rh\_pnc\_nb\_2days)

label var rh\_pnc\_nb\_2days "PNC check within two days for newborn"

//PNC provider for newborn

\*\* This is country specific and could be different for different surveys, please check footnote of the table for this indicator in the final report.

\*Providers of PNC for home deliveries or checks after discharge

recode m72\_1 (0 = 0 "No check") (11 = 1 "Doctor") (12/13 = 2 "Nurse/Midwife") ( 14/15 = 3 "Other skilled provider") (16/90 = 4 "Non-skilled provider") (96 = 5 "Other") ///  
( else = 9 "Don't know or missing") if age<24 & rh\_pnc\_nb\_2days==1, gen(pnc\_nb\_pv\_home)  
replace pnc\_nb\_pv\_home = 0 if rh\_pnc\_nb\_2days==0 & age<24

\*Providers of PNC for facility deliveries

recode m76\_1 (0 = 0 "No check") (11 = 1 "Doctor") (12/13 = 2 "Nurse/Midwife") ( 14/15 = 3 "Other skilled provider") (16/90 = 4 "Non-skilled provider") (96 = 5 "Other") ///  
( else = 9 "Don't know or missing") if age<24 & rh\_pnc\_nb\_2days==1 , gen(pnc\_nb\_pv\_hf)  
replace pnc\_nb\_pv\_hf = 0 if rh\_pnc\_nb\_2days==0 & age<24

\*Combine two PNC provider variables

clonevar rh\_pnc\_nb\_pv = pnc\_nb\_pv\_hf  
replace rh\_pnc\_nb\_pv = pnc\_nb\_pv\_home if (pnc\_nb\_pv\_hf ==9) & rh\_pnc\_nb\_2days ==1 & age<24

\*label variable

label var rh\_pnc\_nb\_pv "Provider for newborns's PNC check"

```
}
}
```

\*survey does not have newborn PNC indicators

```
if m70s_included==0 {
```

```
* replace indicators as missing
```

```
gen rh_pnc_nb_timing = .
```

```
gen rh_pnc_nb_2days = .
```

```
gen rh_pnc_nb_pv = .
```

```
}
```

\*=====TABLES PRODUCED=====

```
ta rh_pnc_nb_2days[iw=w] //for Newborn
```

PNC check within two

days for newborn

	Freq.	Percent	Cum.
--	-------	---------	------

No Visit within 2 days	798.142952	25.19	25.19
------------------------	------------	-------	-------

visit within 2 days	2,370.5574	74.81	100.00
---------------------	------------	-------	--------

Total	3,168.7003	100.00	
-------	------------	--------	--

```
ta rh_pnc_wm_2days[iw=w] // for Mom or mother
```

PNC check

within two

days for

mother

	Freq.	Percent	Cum.
--	-------	---------	------

Not in 2 days	932.12392	29.42	29.42
---------------	-----------	-------	-------

Within 2 days	2,236.5764	70.58	100.00
---------------	------------	-------	--------

Total	3,168.7003	100.00	
-------	------------	--------	--

Subject: Re: Post Natal Care check  
Posted by [Shireen-DHS](#) on Tue, 30 Nov 2021 19:29:26 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hello Francois,

Hope you are doing well.

I was able to match the tables using the GitHub code. I suspect the problem is because you computed age and did not use b19\_01 for age. If you check the RH\_age\_period.do file you will see that if b19 is present in the data file, then this should be used for age. So instead of age=v008- b3\_01 you should use gen age=b19\_01. Then run the entire RH\_PNC.do file to get a match. See the attached Stata output screenshot.

Hope this helps.

Best,

Shireen Assaf

The DHS Program

---

#### File Attachments

1) [RW\\_PNC.png](#), downloaded 205 times

---

---

Subject: Re: Post Natal Care check  
Posted by [Francois](#) on Wed, 01 Dec 2021 11:00:30 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Dear Shireen,

Thank you very much. It works!

So is this the same variable used determining age for ANC while calculating the women "Received 2+ tetanus injections during last pregnancy" and women "Protected against neonatal tetanus"?  
Also, the variable "Period" is similar with which variable?  
Thank you very much

Francois

---

---

Subject: Re: Post Natal Care check  
Posted by [Francois](#) on Wed, 01 Dec 2021 11:20:15 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Dear Shireen,

Ignore the questions for ANC!! Also, it works!!

Thank you

---

---

Subject: Re: Post Natal Care check

Posted by [Shireen-DHS](#) on Wed, 01 Dec 2021 13:27:45 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Great to hear it worked for you.

Yes the age would be the same for the ANC. Please read the main file and the age\_period file. The period is usually births in the last past 5 years for ANC indicators (i.e. 60 months) or the past 2 years for PNC indicators (i.e. 24 months). For Rwanda you dont need to change anything because the code is set up for these periods. For other surveys you need to see what it says in the table title (births in the last 5 years or other).

Best,  
Shireen

---