Subject: Merging Female (IR) file and Male (MR) files Posted by G Kumar S on Fri, 07 Jul 2023 10:44:09 GMT

View Forum Message <> Reply to Message

Hello all,

I am in the in the process of analyzing the Indian DHS data (NFHS-V (2019-21)). While looking for some references, I came across a paper titled 'Knowledge of HIV/AIDS and its determinants in India: Findings from the National Family Health Survey-5 (2019-21)'.

This paper merged the HIV module (respondents aged 15-49 years) female (108785) and male (93267) files of NFHS-5 (2019-21) into a single dataset. The outcome variable of the paper was Knowledge of HIV/AIDS, with gender as one of the independent variables of the analysis.

- 1. Is this okay to combine these files for deriving combined figures?
- 2. If so, what weights are to be used?

Thank you in advance.

The link of the paper is http://www.populationmedicine.eu/Knowledge-of-HIV-AIDS-and-its-determinants-in-India-nFindings-from-the-National-Family, 163113,0,2.html

Subject: Re: Merging Female (IR) file and Male (MR) files Posted by Janet-DHS on Mon, 10 Jul 2023 14:46:09 GMT

View Forum Message <> Reply to Message

Following is a response from DHS staff member, Tom Pullum:

Men were subsampled in NFHS-4 and -5, so there are only about 1/6 as many men as women. If the cases for women and men were put into a single file, and a weight was calculated as v005 for women and 6*mv005 for men, the weighted sample would be approximately representative of the population of women and men combined. This simple calculation of a combined weight is probably not optimal but would be useful. (In regressions these weights would automatically be re-normalized to have a mean of 1.) It would be safest to give separate estimates for women and men separately rather than for the pooled sample.

Subject: Re: Merging Female (IR) file and Male (MR) files Posted by G Kumar S on Wed, 12 Jul 2023 04:29:21 GMT

View Forum Message <> Reply to Message

Thank you very much for the clarification.