
Subject: Calculating perinatal death

Posted by [fahmidarima7](#) on Wed, 13 Nov 2024 07:45:38 GMT

[View Forum Message](#) <> [Reply to Message](#)

****For ENMR, use BR file*******

* create a child Alive or Died variable using the b5 variable

gen alive=b5

lab def alive 0 "Died" 1 "Alive"

lab val alive alive

lab var alive "Alive or Died by the time of survey"

**** Age at death using variables b6 and b5**

gen age_death=.

replace age_death = 0 if b6<=106 & b5==0

replace age_death = 1 if b6>=107 & b6<=130 & b5==0

replace age_death = 2 if (b6> 130 & b6<=999) | b5==1

lab def age_death 0 "Early Neonatal Death" 1 "Late Neonatal Death" 2 "Survived Neonatal Period"

lab val age_death age_death

lab var age_death "Neonatal Mortality Status"

tab age_death

gen wt=v005/1000000

svyset v021 [pw=wt], strata(v023)

svy: tab age_death if v008-b3<60, per count form(%7.3g)

svy: tab age_death if v008-b3<60, per col form(%7.3g)

****For stillbirth, use IR file*****

*Calculate still births in the last 5 years

gen stillbirths = 0

gen births = 0

gen nlbirths = 0

*Set length of calendar to use

gen callen = v018 + 59

* If calendar is aligned right (as in original dataset), use the following:

gen beg = v018

gen end = callen

* If calendar is aligned left (as it appears to be), use the following:

*gen beg = 1

*gen end = 60

* Loop through calendar summing births, non-live pregnancies and stillbirths

forvalues i = 1/80 {

 replace births = births+1 if `i' >= beg & `i' <= end & substr(vcal_1,`i',1) == "B"

 replace nlbirths = nlbirths+1 if `i' >= beg & `i' <= end & substr(vcal_1,`i',1) == "T"

```

    replace stillbirths = stillbirths+1 if `i' >= beg & `i' <= end & substr(vcal_1,`i',7) == "TPPPPP"
}

* total pregnancies in last 5 years
gen totpreg5 = births+nlbirths
* total pregnancies of 7+ months in last 5 years (all live births, plus the stillbirths)
gen totpreg7m = births+stillbirths

* Create weight variable.
gen wgt = v005/1000000

* Set up svyset parameters for complex samples.
svyset v021 [pweight=wgt], strata(v023)

* Produce number of stillbirths
svy: tab stillbirths, cell count
*****
*****Merge BR and IR file*****
* preamble
numlabel,add
set more off
clear
////merge birth recode and individual recode/////
//merge with individual recode
use "/Users/Documents/BDIR81DT/BDIR81FL.DTA"

**BR file**
use "/Users/Documents/BDBR81DT/BDBR81FL.DTA", replace

keep b3 b5 b6 v005 v008 v021 v023

**merge**
merge 1:1 _n using "/Users/Documents/BDIR81DT/BDIR81FL.DTA"

keep if _merge==3
*****

```

I have prepared a syntax for calculating the perinatal mortality rate in Stata with guidance from this forum. However, my results do not match the numbers in the official report, and I am unsure where I might have gone wrong. Could you please review my syntax to help identify any errors? Additionally, I am finding the syntax on GitHub for calculating perinatal deaths quite complex. If possible, could you provide a simplified version of perinatal death equation in Stata?

Subject: Re: Calculating perinatal death
 Posted by [Janet-DHS](#) on Sat, 16 Nov 2024 00:42:37 GMT
[View Forum Message](#) <> [Reply to Message](#)

Following is a response from DHS staff member, Tom Pullum:

With DHS-8 data and the birth histories it is now much easier to calculate the perinatal mortality rate. I will paste below the Stata lines to do it, using the new GR file. It includes code for one covariate (wealth quintiles) and the total.

We have written new Stata code for Chapters 5 and 8 but it may not yet be posted on the GitHub site.

* Table 8.4 in Bangladesh 2022 DHS final report

* Use the GR file for DHS8 surveys

```
use "...BDGR81FL.DTA", clear
```

* Reduce the file to births and stillbirths in months 0-59 before the survey

```
keep if p19<60 & (p32==1 | p32==2)
```

```
gen wt=v005/1000000
```

* Construct the outcome variables

```
gen stillbirth=0
```

```
gen enn_death=0
```

```
replace stillbirth=1 if p32==2
```

```
replace enn_death=1 if p32==1 & p6<=106
```

```
replace stillbirth=stillbirth*wt
```

```
replace enn_death=enn_death*wt
```

```
gen case=wt
```

collapse (sum) stillbirth enn_death case, by(v190)

gen st_rate=1000*stillbirth/case

gen enn_rate=1000*enn_death/case

gen peri_rate=st_rate+enn_rate

gen ratio=stillbirth/enn_death

list, table clean

collapse (sum) stillbirth enn_death case

gen st_rate=1000*stillbirth/case

gen enn_rate=1000*enn_death/case

gen peri_rate=st_rate+enn_rate

gen ratio=stillbirth/enn_death

list, table clean

Subject: Re: Calculating perinatal death

Posted by [fahmidarima7](#) on Wed, 27 Nov 2024 06:22:20 GMT

[View Forum Message](#) <> [Reply to Message](#)

Wow, it's great! I also want to know regarding essential newborn care, if data on "baby dried <5 min of birth" and "baby wrapped <5 min of birth" are available in DHS-8. If so, could you let me know the variable numbers.

Subject: Re: Calculating perinatal death

Posted by [Janet-DHS](#) on Tue, 03 Dec 2024 16:01:46 GMT

[View Forum Message](#) <> [Reply to Message](#)

Following is a response from DHS staff member, Tom Pullum:

Those variables are not included in DHS surveys. To find out which antenatal and postnatal variables are in the data, you can open the GR file in Stata and enter "describe m*". (Also

"describe s*" in other surveys but I see nothing relevant among the "s" variables in this survey.) The variables most relevant to your question have the prefix "mnb" ("nb" for newborn.) You can also look at the questionnaire.
