
Subject: When to use whole numbers versus decimals
Posted by [DHS user](#) on Fri, 15 Feb 2013 14:54:45 GMT
[View Forum Message](#) <> [Reply to Message](#)

When is it acceptable to round to the nearest whole number and when do I need to include one decimal place?

Subject: Re: When to use whole numbers versus decimals
Posted by [Sarah-DHS](#) on Fri, 15 Feb 2013 15:10:00 GMT
[View Forum Message](#) <> [Reply to Message](#)

In general, percentages can be rounded to the nearest whole number. Examples include (but are not limited to): contraceptive prevalence, births delivered in health facilities, children fully immunized, households with an insecticide-treated mosquito net. The exception for rounding percentages is HIV prevalence. Typically, HIV prevalence is presented including one decimal place. However, in some countries where HIV prevalence is higher (mainly southern Africa), HIV prevalence can be rounded to the nearest whole number.

Examples of DHS data that are presented with one decimal place include:

Total Fertility Rate: 6.4 children per woman

Medians: Median Age at First Birth is 18.7 years, Median Age at First Marriage, Median Age at First Sexual Intercourse

Means: Mean Ideal Number of Children is 4.1 children, Average Number of Mosquito Nets per Household, Mean Number of Sexual Partners in Lifetime

Rates and ratios are presented as whole numbers. They should also include the denominator. For example:

Age-specific fertility rate: 116 births per 1,000 women

Under-five mortality rate: 108 deaths per 1,000 live births

Maternal mortality ratio: 454 deaths per 100,000 live births
