
Subject: mean and median duration of breastfeeding
Posted by [Chamberlain](#) on Thu, 02 May 2013 13:56:35 GMT

[View Forum Message](#) <> [Reply to Message](#)

Further, I am writing to request for help to estimate mean and median duration of breastfeeding, postpartum amenorrhoea and insusceptibility, preferably using the lifetable. I would like to examine the patterns by age groups and relate them to contraception adoption. Postpartum variables are not broken down by age and are not readily available in DHS reports, I would appreciate if you could suggest the Stata/spss lifetable commands for computation of the variable or something similar. I have tried to estimate using the Prevalence /Incidence method in SPSS but I am getting estimates which are higher than DHS estimates and some of them implausible for the age group 15-19.

Subject: Re: mean and median duration of breastfeeding
Posted by [Bridgette-DHS](#) on Wed, 22 May 2013 16:39:43 GMT

[View Forum Message](#) <> [Reply to Message](#)

Here is a response from one of our DHS experts Shea Rutstein.

Estimates of duration of the postpartum variables (breastfeeding, amenorrhea, insusceptibility) can be based on recall of durations or current status. While an appropriate methodology for calculating means from recall of durations is using a life table, the well known issues of substantial heaping on durations which are multiples of 3 or 6 indicate that the results will be biased and inaccurate. The DHS reports therefore use current status rather than recall. They also use medians, which are less subject to the effects of long tails than are means. The prevalence-incidence method of calculating a mean is also a current status measure. However, it is based on a stationary population, meaning that there it assumes a constant numeric flow of births. This assumption is violated in many cases, especially for the ends of the fertile age distribution since there were far fewer births to women now 15-19 three years ago (when they were 12-16) than one year ago (when they were 14-18) and so the means will be wrong. The P/I mean is not so bad for the whole age group 15-49 but is somewhat affected by the few women breastfeeding (amenorrheic) for a long time since it is a mean.

I would use an analysis which looks at whether the woman was currently breastfeeding or amenorrheic or insusceptible at the time of the survey and whether she had begun using contraception since the last birth.

Subject: Re: mean and median duration of breastfeeding
Posted by [fllerena](#) on Tue, 16 Jul 2013 22:18:06 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear all. I am Freddy Llerena from Ecuador. I am writing the final report for the last DHS from Ecuador (2012). I have done almost all calculations, but there are a few indicators the I could not

estimate. So, Could you share a syntax (SPSS, STATA, or other), in order to estimate postpartum amenorrhea, insusceptibility and postpartum sexual abstinence. On the other hand, do you have a code for measure infant mortality. Both from DHS data set.

Thanks and kind regards

Subject: Re: mean and median duration of breastfeeding

Posted by [Sarah B](#) on Fri, 19 Jul 2013 14:41:14 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Freddy,

Here's some code I wrote to match the table 6.8 in the Ethiopia 2005 data, which is the table on postpartum amenorrhea, insusceptibility and postpartum sexual abstinence. Please note that this table (and code) uses current status data, rather than recall of durations, which Shea mentions in his response.

** Use the KR data file, which contains all births in the 5 years prior to the survey

use "C:\DATA\ETKR51FL.DTA", clear

** Calculate age of the child in months

g agem=v008-b3

** generate sampling weight

g wgt = v005/1000000

**The denominator for the table = women with children <36m old.

**select only 2nd birth for multiple births -- assigned midx=1

ta b0 if agem<36 & (b0==0 | b0==2) [iw=wgt]

**note that this matches the total denominator.

**amenorrheic

g amen=0 if agem<36 & (b0==0 | b0==2)

*if currently pregnant, make not amenorrheic. Select for numerator only information from most recent birth (m6).

recode amen 0=1 if m6==96 & v213!=1 & midx==1

ta amen [iw=wgt]

**abstaining - same denominator and selections as for amenorrheic

g abst=0 if agem<36 & (b0==0 | b0==2)

recode abst 0=1 if m8==96 & v213!=1 & midx==1

ta abst [iw=wgt]

**insuceptible = amenorrheic or abstaining

g insuc = 0 if agem<36 & (b0==0 | b0==2)

recode insuc 0=1 if abst==1 | amen==1

ta insuc [iw=wgt]

**All match ET table 6.8

I hope this helps with your first question.

Cheers,
Sarah

Subject: Re: mean and median duration of breastfeeding
Posted by [Bridgette-DHS](#) on Mon, 22 Jul 2013 13:48:47 GMT
[View Forum Message](#) <> [Reply to Message](#)

Here is a response from one of our STATA experts Tom Pullum, that should answer your question.

Freddy --The attached programs will calculate the standard mortality rates, and confidence intervals for them, in Stata. The rates agree exactly with the DHS CPro programs. The confidence intervals are close but do not match exactly.

The programs are just text files but you can insert them into the Stata do-file editor. You will have to change the paths and the file name. This is set up for a Nigeria (NG) survey.

Let me know if you have problems. Cheers--Tom

File Attachments

- 1) [child_mortality_24May2013_NG_Pullum_do.txt](#), downloaded 2109 times
 - 2) [child_mortality_ci_25March2013_do.txt](#), downloaded 1670 times
-

Subject: Re: mean and median duration of breastfeeding
Posted by [Lonkila](#) on Sat, 19 Sep 2015 13:58:02 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hello all, I am Lonkila Zan, working on Burkina Faso 2010 dataset

I used the stat code given by Sarah B and I got exactly the same proportion of amenorehic, abstening ans insusceptible like the Burkina 2010 report.

For the mean and the median of the three variables, I used the calculations explained in the Guide to DHS statistics pages 74-75-76.

But I did not get the same numbers like the report.

Can someone explain me how to procced after the code given by Sarah B.

Thank.

Subject: Re: mean and median duration of breastfeeding

Posted by [Sarah B](#) on Wed, 14 Oct 2015 17:27:07 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Lonkila,

When you say "For the mean and the median of the three variables, I used the calculations explained in the Guide to DHS statistics pages 74-75-76.

But I did not get the same numbers like the report," could you please tell us exactly what you are doing, perhaps by pasting your code here, and exactly which #s you are trying to match? I'll try to take a look at your code and see why your results are different than the final report.

Cheers,
Sarah

Subject: Re: mean and median duration of breastfeeding

Posted by [enansubuga](#) on Mon, 22 Oct 2018 08:34:53 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hello,

Kindly asking for the stata code for deriving the mean, median as shown in Table 5.6 for the Uganda 2016 udhs. I have used the code above and the figures match though less by a few decimal points. However, the code stops on deriving the totals and not the mean and median.

Any assistance is highly rendered.
