
Subject: comprehensive knowledge of HIV AIDS
Posted by [cmjacob](#) on Wed, 29 Jul 2015 19:30:36 GMT
[View Forum Message](#) <> [Reply to Message](#)

I am doing a secondary data analysis of Women's empowerment and HIV-KAB in Zambia and my outcome measure is 'comprehensive knowledge (CK) of HIV aids. I intend to use binary logistic regression for the study and have tried to create the CK variable on SPSS. However I am not able to create it as a 1/0 variable. I have found certain other studies that have used this variable as an outcome measure but they have used stata for the same.

I was wondering if it is possible to create a 1/0 variable on SPSS for comprehensive knowledge (I tried using Compute new variable - and gave the IF command to check if answers to the 5 questions about HIV, were answered correctly ; and also recode into new variable - but it did not work.)

If someone has created this variable on SPSS before it would be great if you could share the syntax with me.

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [Liz-DHS](#) on Thu, 06 Aug 2015 14:34:32 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear User,
We have submitted your query to one of our experts. As soon as we have a response, we will post.
Thank you!

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [Trevor-DHS](#) on Thu, 06 Aug 2015 16:15:51 GMT
[View Forum Message](#) <> [Reply to Message](#)

Try the following code:

* A healthy-looking person can have the AIDS virus.

IF (V756 = 1) Healthy = 1.

VARIABLE LABELS Healthy "A healthy-looking person can have the AIDS virus".

VALUE LABELS Healthy 1 "A healthy-looking person can have the AIDS virus".

* The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC].

IF (V754JP = 0) Mosquito = 1.

VARIABLE LABELS Mosquito "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]".

VALUE LABELS Mosquito 1 "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]".

* The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC] .

IF (V823 = 0) SNmeans = 1.

VARIABLE LABELS SNmeans "The AIDS virus cannot be transmitted by supernatural means

[COUNTRY SPECIFIC]".

VALUE LABELS SNmeans 1 "The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC]".

* A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC] .

IF (V754WP = 0) Sfood = 1.

VARIABLE LABELS SFood "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]".

VALUE LABELS SFood 1 "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]".

* The two most common local misconception.

* Need to pick the two most common misconceptions - typically mosquito bites and sharing food - but this can vary from survey to survey.

COUNT MSConcep = mosquito, Sfood (1).

*COUNT MSConcep = mosquito, SNMeans (1).

* A healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions.

IF (Healthy = 1 & MSConcep = 2) MCRreject = 1.

VARIABLE LABELS MCRreject "".

VALUE LABELS MCRreject 1 "Percentage who say that a healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions".

* Percentage with a comprehensive knowledge about AIDS.

IF (V754CP = 1 & V754DP = 1 & Healthy = 1 & MSConcep = 2) knowAIDS = 1.

VARIABLE LABELS knowAIDS "".

VALUE LABELS knowAIDS 1 "Percentage with a comprehensive knowledge about AIDS".

Note that the selection for the two most common mis-conceptions in the middle of the code is survey specific, and will vary from country to country.

Subject: Re: comprehensive knowledge of HIV AIDS

Posted by [CKAllen](#) on Thu, 25 Aug 2016 21:25:48 GMT

[View Forum Message](#) <> [Reply to Message](#)

Does anyone know what the two most common local misconceptions are for Zambia 2007?

On page xxvii of the ZDHS report, it reports 'sharing food' and 'mosquito bites' as the two most common local misconceptions.

On page 192 and 193, it reports 'supernatural means' and 'mosquito bites' as the two most common local misconceptions.

I found similar inconsistencies in 2013 ZDHS report.

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [Liz-DHS](#) on Fri, 26 Aug 2016 18:33:25 GMT
[View Forum Message](#) <> [Reply to Message](#)

A response from Senior Technical Expert, Dr. Kia Reinis:
Quote:
Dear CK Allen,
Thank you for your question.

The misconceptions pertaining to means of transmission that are the two most common: people can get HIV from mosquito bites and people can get HIV from witchcraft or other supernatural means. This is true in both the 2007 and 2013-14, and the data in the chapter tables indicate such.

Footnote 8 in the MDG table in the 2007 report is in error, it should have referred to supernatural means, and not to sharing food.

The text on page 199 of the 2013-14 report is not correct. The excerpt from the report (shown below) that states sharing food is one of the two most common misconceptions is an error, it should have stated "that a person can get HIV via supernatural means" and not "by sharing food".

Thank you very much for bringing this to our attention.
We appreciate your use of DHS data.

File Attachments

1) [ZM2007pg199.jpg](#), downloaded 1193 times

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [CKAllen](#) on Sat, 27 Aug 2016 14:22:33 GMT
[View Forum Message](#) <> [Reply to Message](#)

Thank you, Dr. Reinis for such a quick response. This will clear up which variables to use. Very helpful!

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [enansubuga](#) on Wed, 11 Jan 2017 16:01:11 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Trevor,

Is it possible to have the STATA syntax for comprehensive knowledge. I guess I can better understand the recoding of variables in STATA.

Thank you.

Elizabeth

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [Trevor-DHS](#) on Thu, 04 May 2017 02:37:07 GMT
[View Forum Message](#) <> [Reply to Message](#)

Its a pretty easy conversion of the SPSS code to Stata, but here it is:

* A healthy-looking person can have the AIDS virus.

```
gen healthy = 1 if v756 == 1
```

```
label var healthy "A healthy-looking person can have the AIDS virus"
```

```
label def Healthy 1 "A healthy-looking person can have the AIDS virus"
```

```
label val healthy Healthy
```

* The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC].

```
gen mosquito = 1 if v754jp == 0
```

```
label var mosquito "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]"
```

```
label def Mosquito 1 "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]"
```

```
label val mosquito Mosquito
```

* The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC] .

```
gen snmeans = 1 if v823 == 0
```

```
label var snmeans "The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC]"
```

```
label def SNmeans 1 "The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC]"
```

```
label val snmeans SNmeans
```

* A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC] .

```
gen sfood = 1 if v754wp == 0
```

```
label var sfood "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]"
```

```
label def SFood 1 "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]"
```

```
label val sfood SFood
```

* The two most common local misconception.

* Need to pick the two most common misconceptions - typically mosquito bites and sharing food - but this can vary from survey to survey.

```
gen msconcep = 0
```

```
replace msconcep = 1 if mosquito == 1
```

* if sharing food is more a more common misconception than supernatural means, use the next line

```

replace msconcep = msconcep + 1 if sfood == 1
* if supernatural means is a more common misconception than sharing food, use the next line
instead
*replace msconcep = msconcep + 1 if snmeans == 1

* A healthy-looking person can have the AIDS virus and who reject the two most common local
misconceptions.
gen mcreject = 0
replace mcreject = 1 if healthy == 1 & msconcep == 2
label var mcreject "Percentage who say that a healthy-looking person can have the AIDS virus
and who reject the two most common local misconceptions"
label def MCRreject 1 "Percentage who say that a healthy-looking person can have the AIDS virus
and who reject the two most common local misconceptions"
label val mcreject MCRreject

* Percentage with a comprehensive knowledge about AIDS.
gen knowaids = 0.
replace knowaids = 1 if v754cp == 1 & v754dp == 1 & healthy == 1 & msconcep == 2
label var knowaids "Percentage with a comprehensive knowledge about AIDS"
label def knowAIDS 1 "Percentage with a comprehensive knowledge about AIDS"
label val knowaids knowAIDSFor Zambia 2007 where supernatural means is a more common
misconception than sharing food, comment out the replace command for sfood, and uncomment
the replace command for snmeans instead.

```

Subject: Re: comprehensive knowledge of HIV AIDS
 Posted by [kamal.chaulagain123@gmail](mailto:kamal.chaulagain123@gmail.com) on Mon, 07 Jan 2019 05:37:07 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear All,
 Can I please get the STATA command for creating comprehensive knowledge on HIV/AIDS?
 Thank you in advance.

Subject: Re: comprehensive knowledge of HIV AIDS
 Posted by [Mlue](#) on Mon, 07 Jan 2019 13:47:26 GMT
[View Forum Message](#) <> [Reply to Message](#)

Hello [kamal.chaulagain123@gmail](mailto:kamal.chaulagain123@gmail.com)

Please see the code below... The code is a bit different to the ones shown above.

I assume you are looking for the code that replicates the figures in the Nepal DHS of 2016.

WOMEN

/*

COMPREHENSIVE KNOWLEDGE ABOUT HIV
WOMEN FILE
NEPAL DHS 2016

*/

```
clear all
set more off
set maxvar 9000
set mem 1g
set matsize 800
cd "...
use "NPIR7HFL", clear
set more off
version 14.2
dtaversion "NPIR7HFL.dta"
pwd
```

```
** WEIGHT VARIABLE
gen weight = v005/1000000
```

```
** SURVEY SET
gen psu = v021
gen strata = v023
```

```
svyset psu [pw = weight], strata(strata) vce(linearized)
```

```
// RENAME
```

```
rename v013 age
rename v106 education
rename v190 wealth
rename v025 residence
rename v024 region
rename sdist district
```

```
////////////////////////////////////
```

/*

```
** VARIABLES
```

v756 A healthy looking person can have HIV (women)
v754jp Can get HIV from mosquito bites (women)
v754wp Can get HIV by sharing food with person who has AIDS (women)
v823 Can get HIV by witchcraft or supernatural means (women)
v005 Woman's individual sample weight

mv756 A healthy looking person can have HIV (men)
mv754jp Can get HIV from mosquito bites (men)
mv754wp Can get HIV by sharing food with person who has AIDS (men)
mv823 Can get HIV by witchcraft or supernatural means (men)
mv005 Man's individual sample weigh

*/

cap drop age2
recode v012 (15/24=1 "15-24") (25/29=2 "25-29") (30/39=3 "30-39") (40/49=4 "40-49"), gen(age2)
label var age2 "Current age"
label val age2 age2

** INDICATORS

** 01. A HEALTHY-LOOKING PERSON CAN HAVE THE AIDS VIRUS

recode v756 (1=1 "Yes") (else=0 "No"), gen(healthy_look_HIV)
label var healthy_look_HIV "Knows that a healthy-looking person can have HIV"
label val healthy_look_HIV healthy_look_HIV

svy: tab age2 healthy_look_HIV, count format(%4.0f)
svy: tab age2 healthy_look_HIV, percent format(%4.1f) row

** 02. HIV CANNOT BE TRANSMITTED BY MOSQUITO BITES

recode v754jp (0=1 "Yes") (else=0 "No"), gen(transmit_mosquito_bites)
label var transmit_mosquito_bites "Knows that HIV cannot be transmitted by mosquito bites"
label val transmit_mosquito_bites transmit_mosquito_bites

svy: tab age2 transmit_mosquito_bites, count format(%4.0f)
svy: tab age2 transmit_mosquito_bites, percent format(%4.1f) row

/*

```

** 03. HIV CANNOT BE TRANSMITTED BY SUPERNATURAL MEANS
recode v823 (0=1 "Yes") (else=0 "No"), gen(transmit_witchcraft)
label var transmit_witchcraft "Knows that HIV cannot be transmitted by supernatural means
(witchcraft)"
label val transmit_witchcraft transmit_witchcraft

```

```

svy: tab age2 transmit_witchcraft, count format(%4.0f)
svy: tab age2 transmit_witchcraft, percent format(%4.1f) row
*/

```

```

** 03. HIV CANNOT BE TRANSMITTED BY TOUCHING SOMEONE WHO HAS HIV
recode s1006 (0=1 "Yes") (else=0 "No"), gen(transmit_touching)
label var transmit_touching "Knows that HIV cannot be transmitted by touching someone who has
HIV"
label val transmit_touching transmit_touching

```

```

svy: tab age2 transmit_touching, count format(%4.0f)
svy: tab age2 transmit_touching, percent format(%4.1f) row

```

```

** 04. A PERSON CANNOT BECOME INFECTED BY SHARING FOOD WITH A PERSON WHO
HAS HIV
recode v754wp (0=1 "Yes") (else=0 "No"), gen(sharing_food_HIV)
label var sharing_food_HIV "Knows that a person cannot become infected by sharing food with a
person who has HIV"
label val sharing_food_HIV sharing_food_HIV

```

```

svy: tab age2 sharing_food_HIV, count format(%4.0f)
svy: tab age2 sharing_food_HIV, percent format(%4.1f) row

```

```

** 05. Persons who say that a healthy looking person can have the AIDS virus and who reject the
two most common local misconceptions
/*

```

Two most common local misconceptions: the AIDS virus can be transmitted by mosquito bites and a person can become infected by sharing food with a person who has AIDS.

```

*/

```

```

gen reject_misconceptions = 0
replace reject_misconceptions = 1 if healthy_look_HIV == 1 & (transmit_mosquito_bites == 1 &
sharing_food_HIV == 1)
label define reject_misconceptions 0"No" 1"Yes"
label var reject_misconceptions "A healthy looking person can have the AIDS virus - reject the two

```



```
most common local misconceptions"
label val reject_misconceptions reject_misconceptions
```

```
svy: tab age2 reject_misconceptions, count format(%4.0f)
svy: tab age2 reject_misconceptions, percent format(%4.1f) row
```

```
**
=====
**
**
=====
**
**
=====
**
```

**** 06. HIV PREVENTION METHODS**

```
cap drop prevention_methods
gen prevention_methods = 0
replace prevention_methods = 1 if v754cp == 1 & v754dp == 1
label define prevention_methods 0"No" 1"Yes"
label var prevention_methods "Knows both prevention methods"
label val prevention_methods reject_misconceptions
```

```
svy: tab age2 prevention_methods, count format(%4.0f)
svy: tab age2 prevention_methods, percent format(%4.1f) row
```

**** 07. COMPEHENSIVE KNOWLEDGE ABOUT HIV**

```
/*
Two most common local misconceptions: the AIDS virus can be transmitted by
mosquito bites and a person can become infected by sharing food with a
person who has AIDS.
*/
```

```
cap drop comprehensive_HIV
gen comprehensive_HIV = 0
replace comprehensive_HIV = 1 if (prevention_methods == 1 & healthy_look_HIV == 1) &
(transmit_mosquito_bites == 1 & sharing_food_HIV == 1)
label define comprehensive_HIV 0"No" 1"Yes"
label var comprehensive_HIV ""
label val comprehensive_HIV comprehensive_HIV
```

```
svy: tab age2 comprehensive_HIV, count format(%4.0f)
svy: tab age2 comprehensive_HIV, percent format(%4.1f) row
```

**

```
=====
**

keep if healthy_look_HIV != .

**
=====
**

exit

MEN

/*

COMPREHENSIVE KNOWLEDGE ABOUT HIV
MEN FILE
NEPAL DHS 2016

*/

clear all
set more off
set maxvar 9000
set mem 1g
set matsize 800
cd "..."
use "NPMR7HFL", clear
set more off
version 14.2
dtaversion "NPMR7HFL.dta"
pwd

*****
** WEIGHT VARIABLE
gen weight = mv005/1000000

*****

** SURVEY SET
gen psu = mv021
gen strata = mv023

svyset psu [pw = weight], strata(strata) vce(linearized)
```

// RENAME

rename mv013 age
rename mv106 education
rename mv190 wealth
rename mv025 residence
rename mv024 region
rename smdist district

////////////////////////////////////

/*

** VARIABLES

mv756 A healthy looking person can have HIV (men)
mv754jp Can get HIV from mosquito bites (men)
sm706 can get hiv by touching someone
mv754wp Can get HIV by sharing food with person who has AIDS (men)
mv823 Can get HIV by witchcraft or supernatural means (men)
mv005 Man's individual sample weigh

*/

cap drop age2
recode mv012 (15/24=1 "15-24") (25/29=2 "25-29") (30/39=3 "30-39") (40/49=4 "40-49"),
gen(age2)
label var age2 "Current age"
label val age2 age2

** INDICATORS

** 01. A HEALTHY-LOOKING PERSON CAN HAVE THE AIDS VIRUS
recode mv756 (1=1 "Yes") (else=0 "No"), gen(healthy_look_HIV)
label var healthy_look_HIV "Knows that a healthy-looking person can have HIV"
label val healthy_look_HIV healthy_look_HIV

svy: tab age2 healthy_look_HIV, count format(%4.0f)
svy: tab age2 healthy_look_HIV, percent format(%4.1f) row

** 02. HIV CANNOT BE TRANSMITTED BY MOSQUITO BITES
recode mv754jp (0=1 "Yes") (else=0 "No"), gen(transmit_mosquito_bites)

```
label var transmit_mosquito_bites "Knows that HIV cannot be transmitted by mosquito bites"
label val transmit_mosquito_bites transmit_mosquito_bites
```

```
svy: tab age2 transmit_mosquito_bites, count format(%4.0f)
svy: tab age2 transmit_mosquito_bites, percent format(%4.1f) row
```

```
*****
```

```
/*
** 03. HIV CANNOT BE TRANSMITTED BY SUPERNATURAL MEANS
recode mv823 (0=1 "Yes") (else=0 "No"), gen(transmit_witchcraft)
label var transmit_witchcraft "Knows that HIV cannot be transmitted by supernatural means
(witchcraft)"
label val transmit_witchcraft transmit_witchcraft
```

```
svy: tab age2 transmit_witchcraft, count format(%4.0f)
svy: tab age2 transmit_witchcraft, percent format(%4.1f) row
*/
```

```
*****
```

```
** 03. HIV CANNOT BE TRANSMITTED BY TOUCHING SOMEONE WHO HAS HIV
recode sm706 (0=1 "Yes") (else=0 "No"), gen(transmit_touching)
label var transmit_touching "Knows that HIV cannot be transmitted by touching someone who has
HIV"
label val transmit_touching transmit_touching
```

```
svy: tab age2 transmit_touching, count format(%4.0f)
svy: tab age2 transmit_touching, percent format(%4.1f) row
```

```
*****
```

```
** 04. A PERSON CANNOT BECOME INFECTED BY SHARING FOOD WITH A PERSON WHO
HAS HIV
recode mv754wp (0=1 "Yes") (else=0 "No"), gen(sharing_food_HIV)
label var sharing_food_HIV "Knows that a person cannot become infected by sharing food with a
person who has HIV"
label val sharing_food_HIV sharing_food_HIV
```

```
svy: tab age2 sharing_food_HIV, count format(%4.0f)
svy: tab age2 sharing_food_HIV, percent format(%4.1f) row
```

```
*****
```

```
** 05. Persons who say that a healthy looking person can have the AIDS virus and who reject the
two most common local misconceptions
```

```
/*
Two most common local misconceptions: the AIDS virus can be transmitted by
```

mosquito bites and a person can become infected by sharing food with a person who has AIDS.

```
*/  
  
gen reject_misconceptions = 0  
replace reject_misconceptions = 1 if healthy_look_HIV == 1 & (transmit_mosquito_bites == 1 &  
sharing_food_HIV == 1)  
label define reject_misconceptions 0"No" 1"Yes"  
label var reject_misconceptions "A healthy looking person can have the AIDS virus - reject the two  
most common local misconceptions"  
label val reject_misconceptions reject_misconceptions
```

```
svy: tab age2 reject_misconceptions, count format(%4.0f)  
svy: tab age2 reject_misconceptions, percent format(%4.1f) row
```

```
**  
=====  
**  
**  
=====  
**  
**  
=====  
**
```

**** 06. HIV PREVENTION METHODS**

```
cap drop prevention_methods  
gen prevention_methods = 0  
replace prevention_methods = 1 if mv754cp == 1 & mv754dp == 1  
label define prevention_methods 0"No" 1"Yes"  
label var prevention_methods "Knows both prevention methods"  
label val prevention_methods reject_misconceptions
```

```
svy: tab age2 prevention_methods, count format(%4.0f)  
svy: tab age2 prevention_methods, percent format(%4.1f) row
```

**** 07. COMPEHENSIVE KNOWLEDGE ABOUT HIV**

```
/*  
Two most common local misconceptions: the AIDS virus can be transmitted by  
mosquito bites and a person can become infected by sharing food with a  
person who has AIDS.  
*/
```

```
cap drop comprehensive_HIV  
gen comprehensive_HIV = 0  
replace comprehensive_HIV = 1 if (prevention_methods == 1 & healthy_look_HIV == 1) &
```

```
(transmit_mosquito_bites == 1 & sharing_food_HIV == 1)
label define comprehensive_HIV 0"No" 1"Yes"
label var comprehensive_HIV ""
label val comprehensive_HIV comprehensive_HIV
```

```
svy: tab age2 comprehensive_HIV, count format(%4.0f)
svy: tab age2 comprehensive_HIV, percent format(%4.1f) row
```

```
**
=====
**
```

```
keep if healthy_look_HIV != .
```

```
**
=====
**
```

```
exit
```

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [Trevor-DHS](#) on Mon, 07 Jan 2019 15:32:40 GMT
[View Forum Message](#) <> [Reply to Message](#)

Note that the misconceptions are survey specific. In the example given, the code assumes that the two most common misconceptions are: the AIDS virus can be transmitted by mosquito bites and a person can become infected by sharing food with a person who has AIDS. However, the misconceptions must be verified for each survey, and the two most common are used. See the Guide to DHS Statistics and search for "misconception" to find "Comprehensive Knowledge about HIV (Total and Youth)" for more information on this indicator. Note that some survey specific misconceptions are included in a few surveys too.

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [kamal.chaulagain123@gmail](#) on Mon, 07 Jan 2019 16:51:40 GMT
[View Forum Message](#) <> [Reply to Message](#)

Thank you so much. It helped me lot!!!

Subject: Re: comprehensive knowledge of HIV AIDS
Posted by [Jayanta](#) on Sun, 08 Nov 2020 02:56:42 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Trevor,

I have used the stata syntax that you provided for calculating the comprehensive knowledge of HIV/AIDS for NFHS 2015-16 India IR data but the result seems to be different from the DHS report. Could you please help me with how it has been calculated in the case of NFHS 2015-16 data.

Thank you in advance for your help in this regards.

Regards,
Jayanta
