Subject: comprehensive knowledge of HIV AIDS Posted by cmiacob on Wed, 29 Jul 2015 19:30:36 GMT

View Forum Message <> Reply to Message

I am doing a secondary data analysis of Women's empowerment and HIV-KAB in Zambia and my outcome measure is 'comprehensive knowledge (CK) of HIV aids. I intend to use binary logistic regression for the study and have tried to create the CK variable on SPSS. However I am not able to create it as a 1/0 variable. I have found certain other studies that have used this variable as an outcome measure but they have used stata for the same.

I was wondering if it is possible to create a 1/0 variable on SPSS for comprehensive knowledge (I tried using Compute new variable - and gave the IF command to check if answers to the 5 questions about HIV, were answered correctly; and also recode into new variable - but it did not work.)

If someone has created this variable on SPSS before it would be great if you could share the syntax with me.

Subject: Re: comprehensive knowledge of HIV AIDS Posted by Liz-DHS on Thu, 06 Aug 2015 14:34:32 GMT

View Forum Message <> Reply to Message

Dear User.

We have submitted your query to one of our experts. As soon as we have a response, we will post.

Thank you!

Subject: Re: comprehensive knowledge of HIV AIDS Posted by Trevor-DHS on Thu, 06 Aug 2015 16:15:51 GMT

View Forum Message <> Reply to Message

Try the following code:

* A healthy-looking person can have the AIDS virus.

IF (V756 = 1) Healthy = 1.

VARIABLE LABELS Healthy "A healthy-looking person can have the AIDS virus".

VALUE LABELS Healthy 1 "A healthy-looking person can have the AIDS virus".

* The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]. IF (V754JP = 0) Mosquito = 1.

VARIABLE LABELS Mosquito "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]".

VALUE LABELS Mosquito 1 "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]".

* The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC] . IF (V823 = 0) SNmeans = 1.

VARIABLE LABELS SNmeans "The AIDS virus cannot be transmitted by supernatural means

[COUNTRY SPECIFIC]".

VALUE LABELS SNmeans 1 "The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC]".

* A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC] .

IF (V754WP = 0) Sfood = 1.

VARIABLE LABELS SFood "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]".

VALUE LABELS SFood 1 "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]".

- * The two most common local misconception.
- * Need to pick the two most common misconceptions typically mosquito bites and sharing food but this can vary from survey to survey.

COUNT MSConcep = mosquito, Sfood (1).

- *COUNT MSConcep = mosquito, SNMeans (1).
- * A healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions.

IF (Healthy = 1 & MSConcep = 2) MCReject = 1.

VARIABLE LABELS MCReject "".

VALUE LABELS MCReject 1 "Percentage who say that a healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions".

* Percentage with a comprehensive knowledge about AIDS.

IF (V754CP = 1 & V754DP = 1 & Healthy = 1 & MSConcep = 2) knowAIDS = 1.

VARIABLE LABELS knowAIDS "".

VALUE LABELS knowAIDS 1 "Percentage with a comprehensive knowledge about AIDS". Note that the selection for the two most common mis-conceptions in the middle of the code is survey specific, and will vary from country to country.

Subject: Re: comprehensive knowledge of HIV AIDS Posted by CKAllen on Thu, 25 Aug 2016 21:25:48 GMT

View Forum Message <> Reply to Message

Does anyone know what the two most common local misconceptions are for Zambia 2007? On page xxvii of the ZDHS report, it reports 'sharing food' and 'mosquito bites' as the two most common local misconceptions.

On page 192 and 193, it reports 'supernatural means' and 'mosquito bites' as the two most common local misconceptions.

I found similar inconsistencies in 2013 ZDHS report.

Subject: Re: comprehensive knowledge of HIV AIDS Posted by Liz-DHS on Fri, 26 Aug 2016 18:33:25 GMT

View Forum Message <> Reply to Message

A response from Senior Technical Expert, Dr. Kia Reinis:

Quote:

Dear CK Allen,

Thank you for your question.

The misconceptions pertaining to means of transmission that are the two most common: people can get HIV from mosquito bites and people can get HIV from witchcraft or other supernatural means. This is true in both the 2007 and 2013-14, and the data in the chapter tables indicate such.

Footnote 8 in the MDG table in the 2007 report is in error, it should have referred to supernatural means, and not to sharing food.

The text on page 199 of the 2013-14 report is not correct. The excerpt from the report (shown below) that states sharing food is one of the two most common misconceptions is an error, it should have stated "that a person can get HIV via supernatural means" and not "by sharing food".

Thank you very much for bringing this to our attention. We appreciate your use of DHS data.

File Attachments

1) ZM2007pg199.jpg, downloaded 2001 times

Subject: Re: comprehensive knowledge of HIV AIDS Posted by CKAllen on Sat, 27 Aug 2016 14:22:33 GMT

View Forum Message <> Reply to Message

Thank you, Dr. Reinis for such a quick response. This will clear up which variables to use. Very helpful!

Subject: Re: comprehensive knowledge of HIV AIDS Posted by enansubuga on Wed, 11 Jan 2017 16:01:11 GMT

View Forum Message <> Reply to Message

Dear Trevor,

Is it possible to have the STATA syntax for comprehensive knowledge. I guess I can better understand the recoding of variables in STATA.

Elizabeth

Subject: Re: comprehensive knowledge of HIV AIDS Posted by Trevor-DHS on Thu, 04 May 2017 02:37:07 GMT View Forum Message <> Reply to Message

Its a pretty easy conversion of the SPSS code to Stata, but here it is:

* A healthy-looking person can have the AIDS virus.

gen healthy = 1 if v756 == 1

label var healthy "A healthy-looking person can have the AIDS virus"

label def Healthy 1 "A healthy-looking person can have the AIDS virus"

label val healthy Healthy

* The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]. gen mosquito = 1 if v754jp == 0

label var mosquito "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]"

label def Mosquito 1 "The AIDS virus cannot be transmitted by mosquito bites [COUNTRY SPECIFIC]"

label val mosquito Mosquito

* The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC] . gen snmeans = 1 if v823 == 0

label var snmeans "The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC]"

label def SNmeans 1 "The AIDS virus cannot be transmitted by supernatural means [COUNTRY SPECIFIC]"

label val snmeans SNmeans

* A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC] .

gen sfood = 1 if v754wp == 0

label var sfood "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]"

label def SFood 1 "A person cannot become infected by sharing food with a person who has the AIDS virus [COUNTRY SPECIFIC]"

label val sfood SFood

- * The two most common local misconception.
- * Need to pick the two most common misconceptions typically mosquito bites and sharing food but this can vary from survey to survey.

gen msconcep = 0

replace msconcep = 1 if mosquito == 1

* if sharing food is more a more common misconception than supernatural means, use the next line

replace msconcep = msconcep + 1 if sfood == 1

* A healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions.

gen mcreject = 0

replace mcreject = 1 if healthy == 1 & msconcep == 2

label var mcreject "Percentage who say that a healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions"

label def MCReject 1 "Percentage who say that a healthy-looking person can have the AIDS virus and who reject the two most common local misconceptions" label val mcreject MCReject

* Percentage with a comprehensive knowledge about AIDS. gen knowaids = 0.

replace knowaids = 1 if v754cp == 1 & v754dp == 1 & healthy == 1 & msconcep == 2 label var knowaids "Percentage with a comprehensive knowledge about AIDS" label def knowAIDS 1 "Percentage with a comprehensive knowledge about AIDS" label val knowaids knowAIDSFor Zambia 2007 where supernatural means is a more common misconception than sharing food, comment out the replace command for sfood, and uncomment the replace command for snmeans instead.

Subject: Re: comprehensive knowledge of HIV AIDS Posted by kamal.chaulagain123@gmail on Mon, 07 Jan 2019 05:37:07 GMT View Forum Message <> Reply to Message

Dear All.

Can I please get the STATA command for creating comprehensive knowledge on HIV/AIDS? Thank you in advance.

Subject: Re: comprehensive knowledge of HIV AIDS Posted by Mlue on Mon, 07 Jan 2019 13:47:26 GMT View Forum Message <> Reply to Message

Hello kamal.chaulagain123@gmail

Please see the code below... The code is a bit different to the ones shown above.

I assume you are looking for the code that replicates the figures in the Nepal DHS of 2016.

WOMEN

^{*} if supernatural means is a more common misconception than sharing food, use the next line instead

^{*}replace msconcep = msconcep + 1 if snmeans == 1

```
COMPREHENSIVE KNOWLEDGE ABOUT HIV
WOMEN FILE
NEPAL DHS 2016
```

*/

clear all set more off set maxvar 9000 set mem 1g set matsize 800 cd "..." use "NPIR7HFL", clear set more off version 14.2 dtaversion "NPIR7HFL.dta" pwd

** WEIGHT VARIABLE gen weight = v005/1000000

** SURVEY SET gen psu = v021gen strata = v023

svyset psu [pw = weight], strata(strata) vce(linearized)

// RENAME

rename v013 age rename v106 education rename v190 wealth rename v025 residence rename v024 region rename sdist district

** VARIABLES

```
v756 A healthy looking person can have HIV (women)
v754jp Can get HIV from mosquito bites (women)
v754wp Can get HIV by sharing food with person who has AIDS (women)
v823 Can get HIV by witchcraft or supernatural means (women)
v005 Woman's individual sample weight
**************************
mv756 A healthy looking person can have HIV (men)
mv754jp Can get HIV from mosquito bites (men)
mv754wp Can get HIV by sharing food with person who has AIDS (men)
mv823 Can get HIV by witchcraft or supernatural means (men)
mv005 Man's individual sample weigh
*/
cap drop age2
recode v012 (15/24=1 "15-24") (25/29=2 "25-29") (30/39=3 "30-39") (40/49=4 "40-49"), gen(age2)
label var age2 "Current age"
label val age2 age2
** INDICATORS
** 01. A HEALTHY-LOOKING PERSON CAN HAVE THE AIDS VIRUS
recode v756 (1=1 "Yes") (else=0 "No"), gen(healthy look HIV)
label var healthy look HIV "Knows that a healthy-looking person can have HIV"
label val healthy look HIV healthy look HIV
svy: tab age2 healthy_look_HIV, count format(%4.0f)
svy: tab age2 healthy_look_HIV, percent format(%4.1f) row
******
** 02. HIV CANNOT BE TRANSMITTED BY MOSQUITO BITES
recode v754jp (0=1 "Yes") (else=0 "No"), gen(transmit_mosquito_bites)
label var transmit mosquito bites "Knows that HIV cannot be transmitted by mosquito bites"
label val transmit mosquito bites transmit mosquito bites
svy: tab age2 transmit_mosquito_bites, count format(%4.0f)
svy: tab age2 transmit_mosquito_bites, percent format(%4.1f) row
******
/*
```

```
** 03. HIV CANNOT BE TRANSMITTED BY SUPERNATURAL MEANS
recode v823 (0=1 "Yes") (else=0 "No"), gen(transmit_witchcraft)
label var transmit_witchcraft "Knows that HIV cannot be transmitted by supernatural means
(witchcraft)"
label val transmit_witchcraft transmit_witchcraft
svy: tab age2 transmit witchcraft, count format(%4.0f)
svy: tab age2 transmit_witchcraft, percent format(%4.1f) row
******
** 03. HIV CANNOT BE TRANSMITTED BY TOUCHING SOMEONE WHO HAS HIV
recode s1006 (0=1 "Yes") (else=0 "No"), gen(transmit_touching)
label var transmit_touching "Knows that HIV cannot be transmitted by touching someone who has
HIV"
label val transmit_touching transmit_touching
svy: tab age2 transmit touching, count format(%4.0f)
svy: tab age2 transmit touching, percent format(%4.1f) row
******
** 04. A PERSON CANNOT BECOME INFECTED BY SHARING FOOD WITH A PERSON WHO
HAS HIV
recode v754wp (0=1 "Yes") (else=0 "No"), gen(sharing_food_HIV)
label var sharing_food_HIV "Knows that a person cannot become infected by sharing food with a
person who has HIV"
label val sharing food HIV sharing food HIV
svy: tab age2 sharing food HIV, count format(%4.0f)
svy: tab age2 sharing_food_HIV, percent format(%4.1f) row
******
** 05. Persons who say that a healthy looking person can have the AIDS virus and who reject the
two most common local misconceptions
/*
Two most common local misconceptions: the AIDS virus can be transmitted by
mosquito bites and a person can become infected by sharing food with a
person who has AIDS.
*/
gen reject_misconceptions = 0
replace reject_misconceptions = 1 if healthy_look_HIV == 1 & (transmit_mosquito_bites == 1 &
sharing_food_HIV == 1)
label define reject misconceptions 0"No" 1"Yes"
label var reject misconceptions "A healthy looking person can have the AIDS virus - reject the two
```

```
most common local misconceptions"
label val reject misconceptions reject misconceptions
svy: tab age2 reject_misconceptions, count format(%4.0f)
svy: tab age2 reject_misconceptions, percent format(%4.1f) row
______
** 06. HIV PREVENTION METHODS
cap drop prevention methods
gen prevention methods = 0
replace prevention methods = 1 if v754cp == 1 & v754dp == 1
label define prevention methods 0"No" 1"Yes"
label var prevention methods "Knows both prevention methods"
label val prevention_methods reject_misconceptions
svy: tab age2 prevention_methods, count format(%4.0f)
svy: tab age2 prevention_methods, percent format(%4.1f) row
********
** 07. COMPEHENSIVE KNOWLEDGE ABOUT HIV
Two most common local misconceptions: the AIDS virus can be transmitted by
mosquito bites and a person can become infected by sharing food with a
person who has AIDS.
*/
cap drop comprehensive_HIV
gen comprehensive HIV = 0
replace comprehensive_HIV = 1 if (prevention_methods == 1 & healthy_look_HIV == 1) &
(transmit mosquito bites == 1 & sharing food HIV == 1)
label define comprehensive HIV 0"No" 1"Yes"
label var comprehensive_HIV ""
label val comprehensive_HIV comprehensive_HIV
svy: tab age2 comprehensive_HIV, count format(%4.0f)
svy: tab age2 comprehensive_HIV, percent format(%4.1f) row
```


keep if healthy_look_HIV != . ______ exit MEN COMPREHENSIVE KNOWLEDGE ABOUT HIV MEN FILE NEPAL DHS 2016 */ clear all set more off set maxvar 9000 set mem 1g set matsize 800 cd "..." use "NPMR7HFL", clear set more off version 14.2 dtaversion "NPMR7HFL.dta" pwd ** WEIGHT VARIABLE gen weight = mv005/1000000** SURVEY SET gen psu = mv021 gen strata = mv023 svyset psu [pw = weight], strata(strata) vce(linearized)

// RENAME rename mv013 age rename mv106 education rename mv190 wealth rename mv025 residence rename mv024 region rename smdist district ** VARIABLES mv756 A healthy looking person can have HIV (men) mv754jp Can get HIV from mosquito bites (men) sm706 can get hiv by touching someone mv754wp Can get HIV by sharing food with person who has AIDS (men) mv823 Can get HIV by witchcraft or supernatural means (men) mv005 Man's individual sample weigh */ cap drop age2 recode mv012 (15/24=1 "15-24") (25/29=2 "25-29") (30/39=3 "30-39") (40/49=4 "40-49"), gen(age2) label var age2 "Current age" label val age2 age2 ** INDICATORS ** 01. A HEALTHY-LOOKING PERSON CAN HAVE THE AIDS VIRUS recode mv756 (1=1 "Yes") (else=0 "No"), gen(healthy_look_HIV) label var healthy look HIV "Knows that a healthy-looking person can have HIV" label val healthy_look_HIV healthy_look_HIV svy: tab age2 healthy_look_HIV, count format(%4.0f) svy: tab age2 healthy_look_HIV, percent format(%4.1f) row ****** ** 02. HIV CANNOT BE TRANSMITTED BY MOSQUITO BITES recode mv754jp (0=1 "Yes") (else=0 "No"), gen(transmit mosquito bites)

label var transmit mosquito bites "Knows that HIV cannot be transmitted by mosquito bites" label val transmit mosquito bites transmit mosquito bites svy: tab age2 transmit_mosquito_bites, count format(%4.0f) svy: tab age2 transmit_mosquito_bites, percent format(%4.1f) row ****** ** 03. HIV CANNOT BE TRANSMITTED BY SUPERNATURAL MEANS recode mv823 (0=1 "Yes") (else=0 "No"), gen(transmit_witchcraft) label var transmit witchcraft "Knows that HIV cannot be transmitted by supernatural means (witchcraft)" label val transmit_witchcraft transmit_witchcraft svy: tab age2 transmit_witchcraft, count format(%4.0f) svy: tab age2 transmit witchcraft, percent format(%4.1f) row ******** ** 03. HIV CANNOT BE TRANSMITTED BY TOUCHING SOMEONE WHO HAS HIV recode sm706 (0=1 "Yes") (else=0 "No"), gen(transmit_touching) label var transmit_touching "Knows that HIV cannot be transmitted by touching someone who has HIV" label val transmit_touching transmit_touching svy: tab age2 transmit touching, count format(%4.0f) svy: tab age2 transmit touching, percent format(%4.1f) row ****** ** 04. A PERSON CANNOT BECOME INFECTED BY SHARING FOOD WITH A PERSON WHO HAS HIV recode mv754wp (0=1 "Yes") (else=0 "No"), gen(sharing_food_HIV) label var sharing food HIV "Knows that a person cannot become infected by sharing food with a person who has HIV" label val sharing food HIV sharing food HIV svy: tab age2 sharing food HIV, count format(%4.0f) svy: tab age2 sharing food HIV, percent format(%4.1f) row ****** ** 05. Persons who say that a healthy looking person can have the AIDS virus and who reject the two most common local misconceptions /*

Two most common local misconceptions: the AIDS virus can be transmitted by

```
mosquito bites and a person can become infected by sharing food with a
person who has AIDS.
*/
gen reject_misconceptions = 0
replace reject_misconceptions = 1 if healthy_look_HIV == 1 & (transmit_mosquito_bites == 1 &
sharing food HIV == 1)
label define reject_misconceptions 0"No" 1"Yes"
label var reject misconceptions "A healthy looking person can have the AIDS virus - reject the two
most common local misconceptions"
label val reject_misconceptions reject_misconceptions
svy: tab age2 reject_misconceptions, count format(%4.0f)
svy: tab age2 reject_misconceptions, percent format(%4.1f) row
** 06. HIV PREVENTION METHODS
cap drop prevention_methods
gen prevention methods = 0
replace prevention methods = 1 if mv754cp == 1 & mv754dp == 1
label define prevention methods 0"No" 1"Yes"
label var prevention methods "Knows both prevention methods"
label val prevention_methods reject_misconceptions
svy: tab age2 prevention_methods, count format(%4.0f)
svy: tab age2 prevention_methods, percent format(%4.1f) row
******
** 07. COMPEHENSIVE KNOWLEDGE ABOUT HIV
Two most common local misconceptions: the AIDS virus can be transmitted by
mosquito bites and a person can become infected by sharing food with a
person who has AIDS.
*/
cap drop comprehensive_HIV
gen comprehensive HIV = 0
replace comprehensive HIV = 1 if (prevention methods == 1 & healthy look HIV == 1) &
```



Subject: Re: comprehensive knowledge of HIV AIDS Posted by Trevor-DHS on Mon, 07 Jan 2019 15:32:40 GMT

View Forum Message <> Reply to Message

Note that the misconceptions are survey specific. In the example given, the code assumes that the two most common misconceptions are: the AIDS virus can be transmitted by mosquito bites and a person can become infected by sharing food with a person who has AIDS. However, the misconceptions must be verified for each survey, and the two most common are used. See the Guide to DHS Statistics and search for "misconception" to find "Comprehensive Knowledge about HIV (Total and Youth)" for more information on this indicator. Note that some survey specific misconceptions are included in a few surveys too.

Subject: Re: comprehensive knowledge of HIV AIDS Posted by kamal.chaulagain123@gmail on Mon, 07 Jan 2019 16:51:40 GMT View Forum Message <> Reply to Message

Thank you so much. It helped me lot!!!

Subject: Re: comprehensive knowledge of HIV AIDS Posted by Jayanta on Sun, 08 Nov 2020 02:56:42 GMT

View Forum Message <> Reply to Message

Dear Trevor,

I have used the stata syntax that you provided for calculating the comprehensive knowledge of HIV/AIDS for NFHS 2015-16 India IR data but the result seems to be different from the DHS report. Could you please help me with how it has been calculated in the case of NFHS 2015-16 data.

Thank you in advance for your help in this regards.

Regards, Jayanta