

---

Subject: Malaria Treatment Variable

Posted by [cbdolan](#) on Thu, 26 Jan 2017 21:03:59 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hello,

I have accessed the Malawi 2012 MIS (Individual Recode) and want to create an outcome variable that includes treatment for child fever in the last two weeks.

Before constructing the variable I want to make sure I understand the coding within the dataset.

In the individual recode there are 6 variables (s404a\_1-6) corresponding to days ago fever started, then 6 treatment variables (s406a1-6). Am I correct that this is one women and up to 6 kids which treatment information is collected? For example s404a\_1 is the number of days the fever started and then s406a1 is that same kid and the number of days after the fever began treatment was sought. Following that pattern s404a\_2 goes with s406a2...s404a3 goes with s406a3...

Is there a better way to construct malaria treatment behavior or am I on the right track?

Thanks,

Carrie

---

---

Subject: Re: Malaria Treatment Variable

Posted by [cbdolan](#) on Mon, 30 Jan 2017 15:19:59 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hello,

I found post #8930 and now understand that the MIS is constructed in similar recode files to the standard DHS (ie. IR, PR, KR...). I also now understand the merge is similar and have correctly merged the 2013 PR and KR using the following code:

```
use Z:\Data\DataIn_Original\2_DHS_MIS\MWKR6HDT_MIS\MWKR6HFL.dta, clear
gen hv001=v001
gen hv002=v002
gen hvidx=b16
sort hv001 hv002 hvidx
save Z:\Data\DataIn_Original\0_Scratch\temp.dta, replace
```

```
use Z:\Data\DataIn_Original\2_DHS_MIS\MWPR6HDT_MIS\MWPR6HFL.dta, clear
sort hv001 hv002 hvidx
merge hv001 hv002 hvidx using Z:\Data\DataIn_Original\0_Scratch\temp.dta
tab _merge
keep if _merge==3
drop _merge
```

\*N=2171

No response to this post is necessary.

Thanks!

---