
Subject: Prevention of Malaria(ITP in pregnancy)
Posted by [kambi](#) on Thu, 18 May 2017 16:19:49 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Team ,

I am using dataset of DR Congo (DSH 2013-2014 : CDBR61FL) on the use of ITP in pregnancy in the case of prevention.

I want to know how to create Two dependent variables that include the number of women (15-49) with a live birth in the last two years preceding the survey who:

1) took 2 or more doses of SP (sulfadoxine-Pyrimethamin) and who received at least one doses during an antenatal visit,

2) received the SP during an antenatal visit.

I read some DHS manuals without understanding some things.

I recoded these variables M14 ,m49a,ml2 and i merged them in stata but I can't arrive to find these two variables.How could you do for recoding these outcomes.

Thanks for your help

Fiston

Subject: Re: Prevention of Malaria(ITP in pregnancy)
Posted by [Liz-DHS](#) on Tue, 23 May 2017 14:00:01 GMT
[View Forum Message](#) <> [Reply to Message](#)

A response from malaria expert, Cameron Taylor:

Quote:

Dear user,

Thanks for your question. To begin with you need to be using the IR file (CDIR61FL) instead of the BR file (CDBR61FL). For the IPTp indicator the unit of analysis is women so you need to be using the women's file. You can find more information about DHS datasets at the following videos.

Introduction to DHS datasets: <https://www.youtube.com/watch?v=fzLNQkkvDeI>

Introduction to DHS Data Structure: <https://www.youtube.com/watch?v=BmiSDPebmgc>

Once in the IR file you will then need to consider the following variables in your indicator calculation:

- During pregnancy took SP/fansidar (m49a_1)
- Took 2+ doses (ml1_1)
- Source of antimalarial (ml2_1)
- Had a delivery of a live baby within the last two years (Date of interview- Date of Birth) or (v008-b3_01)

Regards

Cameron

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [kambi](#) on Tue, 23 May 2017 15:14:18 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear Cameron ,

I followed these two videos but I want to know if there will be a difference in the results by using the BR file, instead of IR file because this BR file includes all births related to the women interviewed , What kind of problems am I going to find in this case?

And by calculating v008-b3 to find the number of women with a birth in the last two years, I don't get the same result as in the Table 12.8 (Result=7168) :

```
gen age_new=v008-b3_1
```

```
keep if age_new <24
```

```
tab age_new
```

(Result = 7154) ,a difference of 14 with what is in the table 12.8 of final report,is it correct the way I calculate ?

I will contact you again for some explanations.

Thanks you for your answer Cameron

FISTON

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [Liz-DHS](#) on Fri, 30 Jun 2017 21:04:19 GMT

[View Forum Message](#) <> [Reply to Message](#)

A response from senior malaria expert, Cameron Taylor:

Quote:

Dear Fiston,

To calculate the IPTp indicator you need to be using the Individual Recode (IR) file since the denominator is "Effectif de femmes ayant eu une naissance au cours des deux années avant l'enquête". The IR file has one row for every woman age 15-49 which you are limiting to women who have had a birth in the past two years. As I noted in your previous post you need to make sure you account for:

- During pregnancy took SP/fansidar (m49a_1)
- Took 2+ doses (ml1_1)
- Source of antimalarial (ml2_1)
- Had a delivery of a live baby within the last two years (Date of interview- Date of Birth) or (v008-b3_01)

You can replicate it using the following code for DRC

use "CDIR61FL.DTA", clear

*Weighting

g wgt = v005/1000000

*Establishing birth in past 2 years

g agemnth=.

replace agemnth=v008-b3_01

lab var agemnth "Establishing birth in past 2 years"

*two or more doses sp/fansidar during anc visit

g sptwoanc=0

replace sptwoanc=1 if m49a_1==1 & (ml1_1 >=2) & (ml1_1 <=97) & (ml2_1==1)

*Proportion of women who received two or more doses of IPTp during their last pregnancy in the last two years, at least one dose of which was received as part of an ANC visit**

tab sptwoanc if agemnth < 24 [iweight=wgt]

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [kambi](#) on Mon, 03 Jul 2017 15:30:06 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear Cameron Taylor and Liz ,

Thank you very much for your help.

I'm using now the women recode (IR) file and I saw your answer but why did you put 97 in your syntax when you created "sptwoanc" (you can see the table of ml1_1).

For our code DRC ,I tried theses syntax and I got now 7168 for the proportion of women who received two or more doses of IPTp during their last pregnancy in the last two years but I have a problème to reproduce the descriptive table ,how can I reproduce the result as in the report for example "the descriptive of age" because I don't able to have the correct table of description as in the report (for example "Tab V013")

Thank you a lot

Fiston Kambi

File Attachments

1) [table_ml1_1.docx](#), downloaded 876 times

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [kambi](#) on Mon, 03 Jul 2017 15:46:42 GMT

[View Forum Message](#) <> [Reply to Message](#)

[quote title=kambi wrote on Mon, 03 July 2017 17:30]Dear Cameron Taylor and Liz ,

Thank you very much for your help.

I'm using now the women recode (IR) file and I saw your answer but i want to know why did you put 97 in your syntax when you created "sptwoanc" (you can see the table of ml1_1).

For our code DRC ,I tried theses syntax and I got now 7168 for the proportion of women who received two or more doses of IPTp during their last pregnancy in the last two years but the problème is now to reproduce the descriptive table,how can I reproduce the result as in the report for example "the descriptive of v013" because I reproduce the descriptive table with the old proportion of 18827.It does not take into account the new proportion of women who have had a birth in the last two years.

Thank you a lot

Fiston Kambi

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [kambi](#) on Mon, 03 Jul 2017 15:47:25 GMT

[View Forum Message](#) <> [Reply to Message](#)

[quote title=kambi wrote on Mon, 03 July 2017 17:46][quote title=kambi wrote on Mon, 03 July 2017 17:30]Dear Cameron Taylor and Liz ,

Thank you very much for your help.

I'm using now the women recode (IR) file and I saw your answer but i want to know why did you put 97 in your syntax when you created "sptwoanc" (you can see the table of ml1_1).

For our code DRC ,I tried theses syntax and I got now 7168 for the proportion of women who received two or more doses of IPTp during their last pregnancy in the last two years but the problème is now to reproduce the descriptive table,how can I reproduce the result as in the report for example "the descriptive of v013" because I reproduce the descriptive table with the old proportion of 18827.It does not take into account the new proportion of women who have had a birth in the last two years.

Thank you a lot

Fiston KAMBI

File Attachments

1) [table_ml1_1.docx](#), downloaded 734 times

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [kambi](#) on Mon, 03 Jul 2017 16:19:26 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear Cameron Taylor and Liz ,

Thank you very much for your help.

I'm using now the women recode (IR) file and I saw your answer but i want to know why did you put 97 in your syntax when you created "sptwoanc" (you can see the table of ml1_1).

For our code DRC ,I tried theses syntax and I got now 7168 for the proportion of women who received two or more doses of IPTp during their last pregnancy in the last two years but the problème is now to reproduce the descriptive table,how can I reproduce the result as in the report for example "the descriptive of v013" because I reproduce the descriptive table with the old proportion of 18827.It does not take into account the new proportion of women who have had a birth in the last two years.

Thank you a lot

Fiston kambu

File Attachments

1) [table_ml1_1.docx](#), downloaded 834 times

Subject: Re: Prevention of Malaria(ITP in pregnancy)

Posted by [Liz-DHS](#) on Sun, 01 Oct 2017 04:25:13 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear User,

Do you still need assistance to this post?

Thank you!
