
Subject: weighting child data

Posted by [slolib86](#) on Tue, 19 Dec 2017 21:18:28 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear all,

I going to investigate factors associated with child height (zscore variable). My main concern is about weights. I read in other messages of the forum, that we have to weight the child data (krfile) using the v005 variable. However, the v005 variable comes from the mother data (irfile) where all the data is unique per caseid variable.

I would like to know if there any problem weighting the child data(krfile) since some values of the v005 variable are repeated per caseid.

Thank you so much for your response

Subject: Re: weighting child data

Posted by [Liz-DHS](#) on Thu, 21 Dec 2017 14:59:16 GMT

[View Forum Message](#) <> [Reply to Message](#)

A response from senior sampling expert, Dr. Ruilin Ren:

Quote:

Regarding the weight variable used for analyzing the child weight, it is depending on the background. If you take only the children living with an interviewed mother, it is better to use the variable V005, the child's mother's weight. If you take all children measured in the household, it is better to use the household weight HV005. Regarding the repeated values of the weight variable (either HV005 or V005), note that the weight is a cluster level weight in DHS surveys, all HH in the same cluster share the same weight; all women in the same cluster share the same weight. So this is not a concern.

Thanks

Ruilin

Subject: Re: weighting child data

Posted by [slolib86](#) on Thu, 11 Jan 2018 19:31:13 GMT

[View Forum Message](#) <> [Reply to Message](#)

Thank you for your response Dr. Ruilin Ren

I have two more questions regarding weights. I am pooling DHS surveys , two surveys per country. The country I selected is Perú. I read that before to run a regression model, I should re-normalize the weights.

I proceed like previous post suggestion:

$V005^* = V005 \times (\text{total females age 15-49 in the country at the time of the survey}) / (\text{number of women age 15-49 interviewed in the survey})$

Since my outcome is child weight:

(1) Could I re-normalize the weights using this conversion factor?

$V005^* = V005 \times (\text{total children age 0-5 in the country at the time of the survey}) / (\text{number of children age 0-5 interviewed in the survey})$

(2) is there other way to re-normalize the weight that you could suggest to me?

Thank you so much

Subject: Re: weighting child data
Posted by [Oquendo](#) on Tue, 16 Jan 2018 13:03:06 GMT
[View Forum Message](#) <> [Reply to Message](#)

We're also in the middle of a project in which we need to account for weight renormalization. No one on our team is sure how to do this. If anyone knows the best way to do this, we would appreciate it a lot. Thank you.

Subject: Re: weighting child data
Posted by [Liz-DHS](#) on Tue, 16 Jan 2018 17:31:10 GMT
[View Forum Message](#) <> [Reply to Message](#)

A response from senior sampler, Dr. Ruilin Ren:

Quote:

For Q#1: The answer is no. The "re-normalize" (rather d-normalize in our words) is to reverse the normalization procedure, so it has to use the women population size and sample size. The children's population size and sample size were not used in the weight normalization.

For Q#2: The answer is no. As stated in the question above, "re-normalization" is to remove the normalization factor from the normalized weight, there is no other way. But if you pool different survey data from the same country, you can "re-normalize" the weight using number of households to approximate, this assumes that the average number of women 15-49 per household is constant across surveys.

$V005^* = V005 \times (\text{total number of residential households in the country at the time of the survey}) / (\text{number of households interviewed in the survey})$

Subject: Re: weighting child data
Posted by [Hassen](#) on Mon, 28 May 2018 04:27:02 GMT
[View Forum Message](#) <> [Reply to Message](#)

Thank you all!!

Subject: Re: weighting child data
Posted by [slolib86](#) on Sat, 11 Aug 2018 22:49:12 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear all, I have a question about calculating a mean after pooled two or more surveys.

I read a post before and it said: "You should not do analyses that combine the surveys in the sense of calculating a single mean for two surveys, because such a number would not refer to a well-defined population and is not interpretable".

Could you please share a more detailed answer or comment about why does this analyses would not refer to a well defined population and additionally, this kind of analyses are not interpretable?

thank you so much for your response

https://userforum.dhsprogram.com/index.php?t=msg&goto=10054&&srch=mean+for+two+survey#msg_10054

Subject: Re: weighting child data
Posted by [Liz-DHS](#) on Fri, 24 Aug 2018 15:18:08 GMT
[View Forum Message](#) <> [Reply to Message](#)

A response from Dr. Tom Pullum:

Quote:

Suppose some variable you are interested in had only been collected in, say, a survey with 1000 cases in Peru in 1990 and a survey with 5000 cases in Kenya in 2010. Would you combine them into a single file with 6000 cases and calculate the mean of the variable? Would that mean be interpretable?
