
Subject: Coding of Frequency Variables
Posted by [ebmoran](#) on Tue, 27 Mar 2018 16:47:10 GMT
[View Forum Message](#) <> [Reply to Message](#)

I am working with a merged data set of 53 DHS surveys for sub-Saharan Africa across time and I am looking to clarify how some of the coding was transformed from the survey format into the recode format. For the variables dealing with the frequency of reading a newspaper, watching TV, and listening to the radio, the frequencies are indicated on surveys as: "ALMOST EVERY DAY1
AT LEAST ONCE A WEEK2
LESS THAN ONCE A WEEK3
NOT AT ALL.....4"

but they appear in R as 0-3. Has the scale been shifted down one integer and 0=Almost every day or has the scale been flipped to indicate a more intuitive order 0=Not at all?

Does this logic hold for other variables that have been recoded similarly?

Thanks!

Subject: Re: Coding of Frequency Variables
Posted by [Bridgette-DHS](#) on Tue, 27 Mar 2018 18:14:11 GMT
[View Forum Message](#) <> [Reply to Message](#)

Following is a response from Senior DHS Stata Specialist, Tom Pullum:

You should specify exactly which survey and variable name you are talking about. Usually the codes do not change, but for this one, "never" or not at all is assigned to code 0 rather than 4.
