
Subject: Calculating minimum meal frequency

Posted by ma4g17@soton.ac.uk on Thu, 30 Aug 2018 15:32:16 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hello,

I am doing my dissertation on the risk factors of stunting in Nepal, and I need to use the three variables V469E V469F V469X to calculate the minimum meal frequency but there are a lot of the data missing, can I know how did you calculate it for the report?

Thank you,

Subject: Re: Calculating minimum meal frequency

Posted by [Liz-DHS](#) on Thu, 30 Aug 2018 16:26:58 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear User, to use these variables V469E, V469F and V469X { at least one child born in the last two years and living with respondent }.

CSPro shows partial coding in the recode as follows:

```
{ times gave child tinned/powdered/fresh milk }
```

```
V469E = Q650DN;
```

```
{ times gave child infant formula }
```

```
V469F = Q650EN;
```

```
{ times gave child yogurt }
```

```
V469X = Q650GN;
```

Take a look at the woman's questionnaire and look at the skips to determine where you would encounter data which is not applicable. Missing data (different from Not Applicable) is identified as:

Quote:

9999, 999, 99, 9 This question should have been answered by the respondent, but the questionnaire

contained no information for this variable (missing data). This code is mainly used in paper based surveys.

Take a look at your frequencies to help you in your analysis. Also take a look at The Guide to DHS Statistics

Please look at pages 127-128 for additional information on how missings are taken into account.
