
Subject: using sample weight

Posted by [aliza](#) on Mon, 06 May 2019 09:42:42 GMT

[View Forum Message](#) <> [Reply to Message](#)

hello, do i have to use sample weight if i want to do combine analysis of 5 data points? If yes where can i get the information on using sample weight for Nepal's DHS data?

Subject: Re: using sample weight

Posted by [boyle014](#) on Mon, 24 Jun 2019 16:38:10 GMT

[View Forum Message](#) <> [Reply to Message](#)

Yes, you do have to use sample weights. You might be able to find the numbers you're looking for in StatCompiler.

If you want confidence intervals to see if the data points are statistically significantly different, I would recommend using IPUMS DHS. Begin by selecting all the Nepal samples. Then select the variables that interest you.

Here is an example of code that is comparing a continuous variable (age at first marriage across time). I imagine you would use something similar for your analysis, depending on your specific variable.

```
replace agefrstmar=. if agefrstmar==99.
```

```
svyset, clear
```

```
svyset [pw=perweight], psu(idhspu) strata(idhsstrata)
```

```
* Create a dummy age variable to use to limit analysis to younger women.
```

```
generate age20_35=0
```

```
replace age20_35=1 if age >= 20 & age <= 35
```

```
* Run a regression predicting age at first marriage by survey year (sample) for  
* younger women.
```

```
svy: regress agefrstmar i.sample if age20_35==1
```

```
margins sample
```

```
marginsplot
```

This will produce a figure with data points and confidence intervals over time.
