

---

Subject: Treatment of symptoms of ARI

Posted by [jubayerbiswas](#) on Mon, 23 Dec 2019 07:48:27 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Hi, there

I am looking for antibiotic use for ARI in the Bangladesh DHS 2014 Data and trying to understand reported ( Bangladesh Demographic and Health Survey 2014; page 147 ) percentage of children under age 5 with symptoms of ARI who received antibiotics. I am using BDKR70FL.DTA . For calculation, i am using the following code in STATA

```
gen antibio=0 if ari==1
replace antibio=1 if ari==1 & (s538g==1 | s538h==1 |s538i==1 |s538j==1 |s538k==1 | s538l==1
|s538m==1 )
```

The problem is that my calculated percentage is lower than the reported percentage

(<https://dhsprogram.com/pubs/pdf/FR311/FR311.pdf>)

Is there anyone who help me to clarify this issue

J.Biswas

---

---

Subject: Re: Treatment of symptoms of ARI

Posted by [Liz-DHS](#) on Thu, 27 Feb 2020 22:25:44 GMT

[View Forum Message](#) <> [Reply to Message](#)

---

Dear User, take a look at this link. It contains the DHS Indicators in Stata. You probably want to look at Chapter 10. Also may want to refer to The Guide to DHS Statistics.

---