
Subject: HOUSEHOLD COOKING FUEL AS A RISK FACTOR FOR
RESPIRATORY ILLNESS AMONG UNDER FIVE CHILDREN

Posted by [IBRO](#) on Mon, 02 Mar 2020 19:09:58 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear all,

I wish to analyse relationship between household cooking fuel and ARI symptoms among under 5 children in Nigeria.

I am using NDHS 2018 datasets

Which file should i use please?

Will it be Ok to use the PR (Household members) file?

Will there be need for weighting?

I am using SPSS software

Kind regards

Subject: Re: HOUSEHOLD COOKING FUEL AS A RISK FACTOR FOR
RESPIRATORY ILLNESS AMONG UNDER FIVE CHILDREN

Posted by [Liz-DHS](#) on Tue, 03 Mar 2020 16:36:07 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear User, for an overview on working with DHS data (including which datasets to use for specific analysis) please click here:. In addition you can click here to access SPSS code to produce Demographic and Health Survey Indicators. As always we recommend using The Guide to DHS Statistics and The Standard Recode Manual. Using Datasets for Analysis is also full of useful information.
