
Subject: HOUSEHOLD COOKING FUEL AS A RISK FACTOR FOR
RESPIRATORY ILLNESS AMONG UNDER FIVE CHILDREN

Posted by [Ibraheema](#) on Mon, 02 Mar 2020 19:09:58 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear all,

I wish to analyse relationship between household cooking fuel and ARI symptoms among under 5 children in Nigeria.

I am using NDHS 2018 datasets

Which file should i use please?

Will it be Ok to use the PR (Household members) file?

Will there be need for weighting?

I am using SPSS software

Kind regards

Subject: Re: HOUSEHOLD COOKING FUEL AS A RISK FACTOR FOR
RESPIRATORY ILLNESS AMONG UNDER FIVE CHILDREN

Posted by [Liz-DHS](#) on Tue, 03 Mar 2020 16:36:07 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear User, for an overview on working with DHS data (including which datasets to use for specific analysis) please click here: [In addition you can click here to access SPSS code to produce Demographic and Health Survey Indicators](#). As always we recommend using [The Guide to DHS Statistics](#) and [The Standard Recode Manual](#). [Using Datasets for Analysis](#) is also full of useful information.
