
Subject: Binary Variables Expanded to Six Categories

Posted by amira.elshal.1@city.ac.uk on Sun, 16 Mar 2014 20:55:49 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear Sirs,

I am working on the Individual Recode files of Egypt's EDHS 2000 and 2008 using Stata.

I understand that a variable such as who attended at the delivery (m3) is coded by DHS as a set of binary variables (m3a, m3b, etc.) for each possible option (doctor, nurse/midwife, etc.). What I do NOT understand is why m3a is expanded into m3a_1, m3a_2,...m3a_6. Is that to deal with the case of a woman giving birth to 6 children for example?

Thank you.

Amira

Subject: Re: Binary Variables Expanded to Six Categories

Posted by [Reduced-For\(u\)m](#) on Mon, 17 Mar 2014 18:17:30 GMT

[View Forum Message](#) <> [Reply to Message](#)

Amira,

The _1, _2.... are in fact for different children - all children born within the last 3/5 years. From the recode manual (<https://www.dhsprogram.com/pubs/pdf/DHSG4/Recode4DHS.pdf>)

The maternity history contains up to six entries, relating to births in the three/five years preceding interview. The use of three/five years is country specific. The entries are in reverse order, such that the first entry relates to the last birth in the last three/five years. There is an entry for all children born in the last three/five years including all twins. The period of three/five years includes months 0 to 35/59 prior to the interview, with month 0 being the month of interview. If there are more than six births in the last three/five years then only the last six are included in the maternity history. Each of the following variables, duration of breastfeeding (M5), duration of postpartum amenorrhea (M7) and duration of postpartum abstinence (M9) may have several cases coded 97 "Inconsistent" since the duration of breastfeeding, amenorrhea or abstinence was impossible in the interval between the birth and the following birth or date of interview if the most recent birth (only the date of interview in the case of breastfeeding).

Subject: Re: Binary Variables Expanded to Six Categories

Posted by amira.elshal.1@city.ac.uk on Wed, 19 Mar 2014 13:34:12 GMT

[View Forum Message](#) <> [Reply to Message](#)

Thank you for your informative reply.

The question now is how I can calculate a maternal indicator if a woman has had more than 1 birth during the last five years.

For example, according to the Guide to DHS Statistics, the ANCC (at least four visits) is 100 times the quotient of the numbers of women who received antenatal care for their last birth, according to grouped number of visits (four visits), divided by the number of women with a birth in the last five years, expressed as a percentage. How can I update this formula to account for women having more than 1 birth during the last five years?

Thank you again.

Amira

Subject: Re: Binary Variables Expanded to Six Categories

Posted by [user-rhs](#) on Wed, 19 Mar 2014 16:57:35 GMT

[View Forum Message](#) <> [Reply to Message](#)

Amira,

I would strongly advise you to read the DHS Recode Manual (link: https://www.dhsprogram.com/pubs/pdf/DHSG4/Recode6_DHS_22March2013_DHSG4.pdf) and the questionnaires at the end of the DHS Final Report for Egypt to familiarize yourself with the data. This is what I personally do whenever I'm stumped about what variable to use or the correct way to use a variable I'm interested in. There is a lot of good information in there you can use: what questions were asked, how were they stored in the dataset, what the _1, _2, _3 in a variable name mean, why some variables have missing values for all observations, etc.

The answer to your question is answered in the DHS Recode Manual.

RHS

NB. FYI, typically the antenatal and postnatal care variables are asked only of the most recent birth, instead of ALL of the births in the last 5 years presumably to improve the reliability of the response and minimize recall bias.

Subject: Re: Binary Variables Expanded to Six Categories

Posted by [amira.elshal.1@city.ac.uk](#) on Wed, 18 Jun 2014 16:08:57 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear RHS,

I have taken your advice and checked both the DHS Recode Manual and Egypt's EMW questionnaire. I have also checked the individual data file I have. However, in Egypt, I have noticed that antenatal and postnatal care variables are asked for all births, not only of the most recent birth. So, if 1 woman gave birth to 2 children in the last five years, I know how many ANC visits she had for each of her 2 children. Does I have to account for the ANC visits of her last birth only when I calculate the variable "ANC - at least 4 visits"? Or her ANC visits for both children?

Thanks.

Amira

Subject: Re: Binary Variables Expanded to Six Categories

Posted by [user-rhs](#) on Wed, 18 Jun 2014 17:22:04 GMT

[View Forum Message](#) <> [Reply to Message](#)

It depends on your research question. If you are interested in # ANC visits per pregnancy for all pregnancies, then you would use all of the data from each woman. If you are interested in the most recent pregnancy, then you would use just the variables with subscript _1 in the name. Note that if you use all of a woman's pregnancies, then you would have to account for "nesting" within the individual woman (as her care-seeking behaviour during pregnancy may be the same for all pregnancies).
