
Subject: Calculating the current use of contraceptive by age group

Posted by [sha_016](#) on Tue, 25 Mar 2014 16:29:32 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear Concern

I am analyzing the DHS-2007 data set for Bangladesh to see the current use of contraceptive by age group. A table of this information is already in the BDHS-2007 report. I was trying to compare the results of the report with the results I come up from the data set.

I found dissimilarity between BDHS-2007 report with my results. I am not sure whether I made the correct commands or not?

Here are the commands I followed:

```
gen strataid = v022
```

```
gen psu = v021
```

```
gen weight = v005/1000000
```

```
svyset psu (pw=weight), strata (strataid)
```

```
gen agegr= v012
```

```
recode agegr (15/19=1) (20/24=2) (25/29=3) (30/34=4) (35/39=5) (40/44=6) (45/49=7)
```

```
svy linearized, subpop(if v213==0) : proportion v312, stdize(weight) stdweight(v005) over (agegr)
```

After running these commands I got a table and results of this table is not the same with the BDHS-2007 report.

Could you please tell me about my possible errors or the correct ways of doing this?

Looking forward to hearing from you,

Thank you,

Shahab

Subject: Re: Calculating the current use of contraceptive by age group

Posted by [Liz-DHS](#) on Tue, 25 Mar 2014 16:46:09 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear User,

Can you please provide the table number and the page number of the table you are looking at in the final report of the Bangladesh DHS 2007?

Thank you!

Subject: Re: Calculating the current use of contraceptive by age group

Posted by [sha_016](#) on Tue, 25 Mar 2014 17:30:00 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear

Table number is 5.5 on the page 60 in the report of DHS-2007 Bangladesh.

Thank you,

Shahab

Subject: Re: Calculating the current use of contraceptive by age group

Posted by [Liz-DHS](#) on Tue, 25 Mar 2014 19:49:11 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear USer,

I am not a programmer, but here is some code in CSPRo that may help in figuring things out.

The row variables are:

V013w age, value labels 1 15-19 2 20-24 3 25-29 4 30-34 5 35-39 6 40-44 7 45-49 8 50-54
(the other row variables in the table definition are other background characteristics which do not display in this particular table)

The Column variables:

useany value label 1 any method
usemod value label 1 any modern method
modmet2 Modern method, value labels 3 Pill 4 IUD 5 Injectables 6 Implants 7 Male condom 1
Femlae sterilization 2 Male sterilization
8 Female condom 9 Diaphragm 10 Foam/Jelly 11 LAM 12 Emergency
contraception
usetrad Value label 1 any traditional method
tradmet2 Traditional method, value labels 13 Periodic abstinence 14 Withdrawal 15 Other
nousing Value label 1 not currently using
totnum2 Value labels 0 Total 1 Number of women

***** *****

Table 5.5 Current use of contraception by age

```
title( "Table 5.5 Current use of contraception by age", " ",  
      "Percent distribution of currently married women age 15-49 by contraceptive",  
      "method currently used and age, Bangladesh 2007" )  
stub( "Age" );
```

```
crosstab float(1) t505 v013w+v102w+v101w+v149w+v218w2+v190w+total  
          useany+usemod+modmet2+usetrad+tradmet2+nousing+totnum2  
exclude(rowzero,colzero,percents,totals,specval)
```

```
{ Table 5.5 processing }  
jtot = tblcol( t505 );  
jmax = jtot - 2;  
do j = 0 while j <= jmax by 1  
  t505[*,j] = t505[*,j] * 100 / t505[*,jtot];  
enddo;  
t505[*,jmax+1] = tblsum( column t505[*, 0:jmax] );  
any = tblcol( t505, useany );  
nouse = tblcol( t505, nousing );  
mod = tblcol( t505, usemod );  
trad = tblcol( t505, usetrad );  
{ total using any method }
```

```

t505[* , any] = tblsum( column t505[* , mod+1:nouse-1] );
{ total using any modern method }
t505[* , mod] = tblsum( column t505[* , mod+1:trad-1] );
{ total using any traditional or other method }
t505[* ,trad] = tblsum( column t505[* ,trad+1:nouse-1] );

```

```
{ table 5.4, 5.5 }
```

```

modmet2 = notappl;
tradmet2 = notappl;
nousing = notappl;
meth = V312;
box meth => method;
  0 => 0;
  1-3 => meth+2; { 3-Pill, 4-IUD, 5-Injections }
  4 => 9;    { 9-Diaphragm }
  5 => 7;    { 7-Male condom }
  6-7 => meth-5; { 1-Female ster, 2-Male ster }
  8,9 => meth+5; { 13-Periodic abstinence, 14-Withdrawal }
  10 => 15;   { 15-Other }
  11 => 6;    { 6-Implants }
  13 => 11;   { 11-Lactational amenorrhea }
  14 => 8;    { 8-Female condom }
  15 => 10;   { 10-Foam/jelly/suppository }
  16 => 12;   { 12-Emergency contraception }
  => default;
endbox;
if V313 = 0 then          { not using }
  nousing = 1
elseif V313 = 3 then     { using modern }
  modmet2 = method;
elseif V313 = 1 | V313 = 2 then { using traditional & others }
  tradmet2 = method;
endif;

if marrwom then

  xtab( t504, rweight ); { smg: moving here june 7, as we only want to show CM women }

  xtab( t505, rweight );

  { smg: this code/table added 8-19-08 }

  if modmet2=7 then { using condom }
    rowMDG5=1; xtab (tMDG5, rweight);
  endif;

  if !(method in 0,default,notappl) then { she's using a method }

```

```
rowMDG5=2; xtab (tMDG5, rweight);  
endif;  
  
endif;
```

Subject: Re: Calculating the current use of contraceptive by age group
Posted by [sha_016](#) on Wed, 26 Mar 2014 08:15:36 GMT
[View Forum Message](#) <> [Reply to Message](#)

Dear Sir/Madam

I am using STATA for analysis. I am not aware of CSPRo.
Thank you for you feedback,
Regards,
Shahab
